

A CRITERION FOR THE LOGARITHMIC DIFFERENTIAL OPERATORS TO BE GENERATED BY VECTOR FIELDS

MATHIAS SCHULZE

ABSTRACT. We study divisors in a complex manifold in view of the property that the algebra of logarithmic differential operators along the divisor is generated by logarithmic vector fields. We give

- a sufficient criterion for the property,
- a simple proof of F.J. Calderón–Moreno’s theorem that free divisors have the property,
- a proof that divisors in dimension 3 with only isolated quasi–homogeneous singularities have the property,
- an example of a non–free divisor with non–isolated singularity having the property,
- an example of a divisor not having the property, and
- an algorithm to compute the V –filtration along a divisor up to a given order.

CONTENTS

1. Logarithmic comparison theorem for free divisors	1
2. V –filtration along subvarieties and divisors	2
3. Symmetric algebra of logarithmic vector fields	5
4. Depth and torsion of symmetric algebras	8
5. Example of generic hyperplane arrangements	9
References	10

1. LOGARITHMIC COMPARISON THEOREM FOR FREE DIVISORS

Let X be a complex manifold of dimension $n \geq 2$, \mathcal{O} the ring holomorphic functions on X , and Ω^\bullet the complex of holomorphic differential forms. Grothendieck’s Comparison Theorem states that the De Rham system \mathcal{O} is regular [Meb89, Thm. 2.3.4]. This is equivalent to the fact that, for any divisor $D \subset X$, the natural morphism

$$\Omega^\bullet(*D) = \mathrm{DR}(\mathcal{O}(*D)) \longrightarrow \mathrm{R}i_*i^{-1}\mathrm{DR}(\mathcal{O}) = \mathrm{R}i_*\mathcal{C}_U,$$

where i is the inclusion $U = X \setminus D \subset X$, is a quasi–isomorphism. Let $\Omega^\bullet(\log D) \subset \Omega^\bullet(*D)$ be the subcomplex of logarithmic differential forms along D [Sai80, Def. 1.2]. The above statement raises the question whether the inclusion $\Omega^\bullet(\log D) \subset \Omega^\bullet(*D)$

1991 *Mathematics Subject Classification.* 32C38, 13A30.

Key words and phrases. free divisor, hyperplane arrangement, logarithmic differential operator, symmetric algebra, V –filtration.

The author is grateful to M. Granger for many valuable discussions and comments and to F.J. Castro–Jiménez, L. Narváez–Macarro, and J.M. Ucha–Enríquez for explaining their results and ideas.

is also a quasi-isomorphism. That is: Can one compute the cohomology of the complement of D by logarithmic differential forms along D ? This turns out to be a property of D called the logarithmic comparison theorem or simply LCT. It is an open problem to characterize the divisors for which LCT holds.

Let $\Theta = \text{Der}_{\mathbb{C}}(\mathcal{O})$ be the \mathcal{O} -module of holomorphic vector fields on X and $\text{Der}(\log D) \subset \Theta$ be the \mathcal{O} -submodule of logarithmic differential operators along D [Sai80, Def. 1.4]. A divisor D is called free if $\text{Der}(\log D)$ is a locally free \mathcal{O} -module. Let \mathcal{D} be the \mathcal{O} -algebra of differential operators on X with holomorphic coefficients and let F be the increasing filtration on \mathcal{D} by the order of differential operators. Let \mathcal{V}^D be the V -filtration along D on \mathcal{D} as defined in Section 2 such that $\mathcal{V}_0^D = \mathcal{D}(\log D)$ is the \mathcal{O} -algebra of logarithmic differential operators along D . F.J. Calderón–Moreno [CM99, Thm. 1] proves that, for a free divisor D , \mathcal{V}_0^D is generated by vector fields, that is $\mathcal{V}_0^D = \mathcal{O}[\text{Der}(\log D)]$. Let S_D be the decreasing filtration on \mathcal{D} which is locally defined by $S_D^k = f^k \cdot \mathcal{D}$ where $f \in \mathcal{O}$ such that $D = (f)$. By Corollary 3, the induced filtration S_D on \mathcal{V}_0^D defined by $S_D^k \mathcal{V}_0^D = \mathcal{V}_0^D \cap (f^k \cdot \mathcal{D})$ reflects the embeddings $\mathcal{V}_k^D \subset \mathcal{D}$. If $\mathcal{V}_0^D = \mathcal{O}[\text{Der}(\log D)]$ then (\mathcal{V}_0^D, S_D) is a filtered $(\mathcal{V}_0^D, \mathcal{V}^D)$ -module.

F.J. Calderón–Moreno and L. Narváez–Macarro [CMNM05, Cor. 4.2] prove that LCT holds for a free divisor D if and only if the complex

$$\mathcal{D} \otimes_{\mathcal{D}(\log D)}^{\mathbb{L}} \mathcal{O}(D) = \mathcal{D} \otimes_{\mathcal{D}(\log D)} \text{Sp}_{\mathcal{D}(\log D)}^{\bullet}(\mathcal{O}(D))$$

is concentrated in degree 0 and the natural multiplication morphism

$$\mathcal{D} \otimes_{\mathcal{D}(\log D)} \mathcal{O}(D) \xrightarrow{\epsilon_D} \mathcal{O}(*D)$$

is an isomorphism. Injectivity of ϵ_D is locally equivalent to $\text{Ann}_{\mathcal{D}(\frac{1}{f})}$ being generated by operators of order 1 where $f \in \mathcal{O}$ such that $D = (f)$. For any divisor D , T. Torrelli proves that the latter condition already implies surjectivity of ϵ_D [Tor04, Prop. 1.3] and conjectures that it is even equivalent to LCT [Tor04, Conj. 1.11]. A problem to verify this conjecture for a free divisor D consists in $\mathcal{D} \otimes_{\mathcal{D}(\log D)} \text{Sp}_{\mathcal{D}(\log D)}^{\bullet}(\mathcal{O}(D))$ not being F -strict in general [CM99, Rem. 4.2.4]. So grading by F does not reduce the problem to a commutative one. But both properties of D in question can be characterized in terms of S_D -strictness: On the one hand, exactness of $\mathcal{D} \otimes_{\mathcal{D}(\log D)} \text{Sp}_{\mathcal{D}(\log D)}^{\bullet}(\mathcal{O}(D))$ in degree k is equivalent to S_D -strictness of the differential of $\text{Sp}_{\mathcal{D}(\log D)}^{\bullet}(\mathcal{O}(D))$ from degree $k-1$ to degree k . On the other hand, injectivity of ϵ_D is equivalent to S_D -strictness of the last differential of $\mathcal{D} \otimes_{\mathcal{D}(\log D)} \text{Sp}_{\mathcal{D}(\log D)}^{\bullet}(\mathcal{O}(D))$.

A solution of the LCT problem seems to require a deeper understanding of the V -filtration in general. There are many questions:

- What are properties of the \mathcal{V}_k^D ?
- When is \mathcal{V}_0^D generated by vector fields?
- When is \mathcal{V}_0^D locally finitely generated?
- What are properties of the embeddings $\mathcal{V}_k^D \subset \mathcal{D}$?

We shall approach the first two questions in this article.

2. V-FILTRATION ALONG SUBVARIETIES AND DIVISORS

Let $Y \subset X$ be a subvariety in X and let $\mathcal{I} \subset \mathcal{O}$ be its ideal. The V-filtration \mathcal{V}^Y along Y is the increasing filtration on \mathcal{D} defined by

$$\mathcal{V}_k^Y = \{P \in \mathcal{D} \mid \forall l \in \mathbb{Z} : P(\mathcal{I}^l) \subset \mathcal{I}^{l-k}\}$$

for all $k \in \mathbb{Z}$. We shall omit the index Y if it is clear from the context. Clearly $\mathcal{V}_k \cdot \mathcal{V}_l \subset \mathcal{V}_{k+l}$ for all $k, l \in \mathbb{Z}$. Hence \mathcal{V}_0 is an \mathcal{O} -algebra and \mathcal{V}_k is an \mathcal{V}_0 -module for all $k \in \mathbb{Z}$.

Example 1. Let $x_1, \dots, x_m, y_1, \dots, y_n$ be coordinates on $X = \mathbb{C}^{m+n}$.

(1) For the submanifold $Y = \{y = 0\}$,

$$\mathcal{V}_k^Y = \{P = \sum_{j_1 - i_1 + \dots + j_n - i_n \leq k} P_{i,j}(x, \partial_x) y_1^{i_1} \partial_{y_1}^{j_1} \dots y_n^{i_n} \partial_{y_n}^{j_n} \in \mathcal{D}\}.$$

(2) For the normal crossing divisor $D = (y_1 \cdots y_n)$,

$$\mathcal{V}_k^D = \{P = \sum_{j_1 - i_1, \dots, j_n - i_n \leq k} P_{i,j}(x, \partial_x) y_1^{i_1} \partial_{y_1}^{j_1} \dots y_n^{i_n} \partial_{y_n}^{j_n} \in \mathcal{D}\}.$$

Denote the complement of the singularities of Y by

$$U_Y = X \setminus \text{Sing}(Y) \xrightarrow{i_Y} X.$$

We shall omit the index Y if it is clear from the context. The V-filtration along a divisor has a special property.

Proposition 2. *Let $D \subset X$ be a divisor. Then $\mathcal{V}^D = (i_D)_* i_D^{-1} \mathcal{V}^D$.*

Proof. We may assume that $D = (f)$ for some $f \in \mathcal{O}$ by the local nature of the statement. Since $\mathcal{V}_k \subset \mathcal{D}$ and \mathcal{D} is a locally free \mathcal{O} -module,

$$i_* i^{-1} \mathcal{V}_k \subset i_* i^{-1} \mathcal{D} = \mathcal{D}.$$

Since $\mathcal{O} \cdot f^{l-k}$ is a free \mathcal{O} -module, $P \in i_* i^{-1} \mathcal{V}_k$ implies

$$P(g \cdot f^l) \in i_* i^{-1} (\mathcal{O} \cdot f^{l-k}) = \mathcal{O} \cdot f^{l-k}$$

for all $g \in \mathcal{O}$ and $l \in \mathbb{Z}$ and hence $P \in \mathcal{V}_k$. □

Corollary 3. *Let $D = (f) \subset X$ with $f \in \mathcal{O}$ be a divisor. Then*

$$\mathcal{V}_k = \begin{cases} f^{-k} \mathcal{V}_0, & k \leq 0, \\ f^{-k} (\mathcal{V}_0 \cap f^k \mathcal{D}), & k \geq 1. \end{cases}$$

Proof. The equalities in question hold on U_D by Example 1 (2) and hence on X by Proposition 2. □

Denote the symbol map for F by

$$\mathcal{D} \xrightarrow{\sigma} \text{gr}^F \mathcal{D}.$$

The decomposition $F_1 \mathcal{D} = \mathcal{O} \oplus \Theta$ defines the \mathcal{O} -module $\text{Der}(\log Y) \subset \Theta$ of logarithmic vector fields along Y by

$$F_1 \mathcal{V}_0 = \mathcal{O} \oplus \text{Der}(\log Y).$$

This definition simplifies to

$$\text{Der}(\log Y) = \{\theta \in \Theta \mid \theta(\mathcal{I}) \subset \mathcal{I}\}$$

by the Leibniz rule and implies involutivity of $\text{Der}(\log Y)$, that is

$$[\text{Der}(\log Y), \text{Der}(\log Y)] \subset \text{Der}(\log Y).$$

Example 4. Let $x_1, \dots, x_m, y_1, \dots, y_n$ be coordinates on $X = \mathbb{C}^{m+n}$.

(1) For the submanifold $Y = \{y = 0\}$,

$$\text{Der}(\log D) = \mathcal{O}\langle \partial_{x_1}, \dots, \partial_{x_m} \rangle + \mathcal{O}\langle y_i \partial_{y_j} \mid 1 \leq i, j \leq n \rangle.$$

(2) For the normal crossing divisor $D = (y_1 \cdots y_n)$,

$$\text{Der}(\log D) = \mathcal{O}\langle \partial_{x_1}, \dots, \partial_{x_m}, y_1 \partial_{y_1}, \dots, y_n \partial_{y_n} \rangle.$$

Let $\mathcal{O}[\text{Der}(\log Y)] \subset \mathcal{D}$ be the image of the tensor algebra

$$\text{T}_{\mathbb{C}} \text{Der}(\log Y) \xrightarrow{\gamma_Y} \mathcal{D}.$$

Then at least $\mathcal{O}[\text{Der}(\log Y)] \subset \mathcal{V}_0^Y$.

Corollary 5. *Let $D \subset X$ be a divisor. Then $\mathcal{V}_0^D = \mathcal{O}[\text{Der}(\log D)]$ if and only if $\mathcal{O}[\text{Der}(\log D)] = (i_D)_* i_D^{-1} \mathcal{O}[\text{Der}(\log D)]$.*

Proof. By Examples 1 and 4, $\mathcal{V}_0^D = \mathcal{O}[\text{Der}(\log D)]$ on U_D . Hence the claim follows from Proposition 2. \square

A divisor $D \subset X$ is called free if $\text{Der}(\log D)$ is a locally free \mathcal{O} -module. By K. Saito [Sai80, Cor. 1.7], $\text{Der}(\log D)$ is reflexive and hence all divisors in dimension $n = 2$ are free. By Example 4 (2), normal crossing divisors are free. In particular, any divisor D is free on U_D .

F.J. Calderón–Moreno [CM99, Thm. 1] proves that $\mathcal{V}_0^D = \mathcal{O}[\text{Der}(\log D)]$ for a free divisor. We give a simple proof of this result.

Corollary 6. *Let $D \subset X$ be a free divisor. Then $\mathcal{V}_0^D = \mathcal{O}[\text{Der}(\log D)]$.*

Proof. By Lemma 7 and grading by F , $\mathcal{O}[\text{Der}(\log D)]$ is a locally free \mathcal{O} -module and hence Corollary 5 applies. \square

Lemma 7. *Let R be a domain and let $P_1, \dots, P_n \in R \cdot T_1 \oplus \cdots \oplus R \cdot T_n = R^n$ be R -linearly independent. Then $R[P_1, \dots, P_n] \subset R[T_1, \dots, T_n]$ is a polynomial ring.*

Proof. Write $P_i = \sum_j p_{i,j} T_j$ with $p_{i,j} \in R$. Then by assumption $p = \det(p_{i,j}) \neq 0$ and hence $R_p[P_1, \dots, P_n]$ is a polynomial ring. Since R is a domain, $R \longrightarrow R_p$ is injective and hence $R[P_1, \dots, P_n]$ is a polynomial ring. \square

In general it is not clear if, or under which conditions, \mathcal{V}_0^Y is a locally finite \mathcal{O} -algebra. Even to compute $F_k \mathcal{V}_0^Y$ is a problem since the definition involves infinitely many conditions. The following result allows one to compute $F_k \mathcal{V}_0^D$ algorithmically.

Proposition 8. *Let x_1, \dots, x_n be coordinates on $X = \mathbb{C}^n$. Let $D = (f) \subset X$ with $f \in \mathcal{O}$ be a divisor. Then, for $P \in F_d \mathcal{D}$, $P \in \mathcal{V}_k^D$ if and only if*

$$(1) \quad \forall \alpha \in \mathbb{N}^n, l \in \mathbb{N} : |\alpha| + l \leq d \Rightarrow P(x^\alpha f^l) \in \mathcal{O} \cdot f^{l-k}.$$

Proof. Let $0 \neq P \in F_d \mathcal{D}$ and assume that condition (1) holds. For $l \in \mathbb{N}$, the vector space $\mathbb{C}[x_1, \dots, x_n]_{\leq d-l}$ is invariant under $x \mapsto Ax + a$ for $a \in \mathbb{C}^n$ and $A \in \text{GL}_n(\mathbb{C})$. Hence, at a smooth point y of D , condition (1) holds for coordinates x_1, \dots, x_n at

y such that $\partial_{x_n}(f)(y) \neq 0$. Then $y_1, \dots, y_{n-1}, t = x_1, \dots, x_{n-1}, f$ are coordinates at y such that

$$\forall \beta \in \mathbb{N}^{n-1}, l \in \mathbb{N} : |\beta| + l \leq d \Rightarrow P_y(y^\beta t^l) \in \mathcal{O}_y \cdot t^{l-k}.$$

Write $P_y = \sum_{|\beta|+l \leq d} p_{\beta,l} \partial_y^\beta \partial_t^l$ with $p_{\beta,l} \in \mathcal{O}_y$ and choose $\gamma \in \mathbb{N}^{n-1}$ and $m \in \mathbb{N}$ such that $|\gamma| + m$ is minimal with $p_{\gamma,m} \neq 0$. Then

$$\gamma! m! p_{\gamma,m} = P(y^\gamma t^m) \in \mathcal{O}_y \cdot t^{m-k}$$

and hence $p_{\gamma,m} \partial_y^\gamma \partial_t^m \in \mathcal{V}_{k,y}$ by Example 1 (2). By increasing induction on $|\gamma| + m$, this implies $P_y \in \mathcal{V}_{k,y}$ for all $y \in U_D$ and hence $P \in \mathcal{V}_k$ by Proposition 2. \square

Example 9. Let x, y, z be coordinates on \mathbb{C}^3 and

$$f = xyz(x + y + z)(x + 2y + 3z).$$

Then $D = (f) \subset \mathbb{C}^3$ is a central generic hyperplane arrangement. Let

$$Q = (x + y + z)(x + 2y + 3z)(3zy^2 \partial_y^2 + (x + 4y - 3z)yz \partial_y \partial_z - 4yz^2 \partial_z^2).$$

Then $Q \in F_2 \mathcal{V}_0^D$ by a SINGULAR [GPS05] computation using Proposition 8. We shall see in Example 13 that $Q \notin F_2 \mathcal{O}[\text{Der}(\log D)]$.

There is another special property of the V-filtration along a divisor.

Proposition 10. *Let $D \subset X$ be a divisor. Then $\text{depth}_x(\mathcal{V}_k^D) \geq 2$ for all $x \in X$ and $k \in \mathbb{Z}$.*

Proof. Let $x \in X$ and $D_x = (f)$ with $f \in \mathcal{O}_x$. Since \mathcal{O}_x is torsion free and $\text{depth}(\mathcal{O}_x) \geq 2$, there is an \mathcal{O}_x -sequence $a_1, a_2 \in \mathfrak{m}_x$ such that a_1 is different from all irreducible factors of f . Let $P \in \mathcal{V}_{k,x}$ with $a_2 \cdot P \in a_1 \cdot \mathcal{V}_{k,x} \subset a_1 \cdot \mathcal{D}_x$. Then $P \in a_1 \cdot \mathcal{D}_x$ since \mathcal{D}_x is a free \mathcal{O}_x -module. But $P(g \cdot f^l) \in \mathcal{O}_x \cdot f^{l-k}$ implies $(a_1^{-1} \cdot P)(g \cdot f^l) \in \mathcal{O}_x \cdot f^{l-k}$ by the choice of a_1 for all $g \in \mathcal{O}$ and $l \in \mathbb{Z}$ and hence $P \in a_1 \cdot \mathcal{V}_{0,x}$. Then $a_1, a_2 \in \mathfrak{m}_x$ is a $\mathcal{V}_{k,x}$ -sequence and hence $\text{depth}_x(\mathcal{V}_k) \geq 2$. \square

3. SYMMETRIC ALGEBRA OF LOGARITHMIC VECTOR FIELDS

The condition in Corollary 5 is difficult to verify in general. Therefore we focus on a case in which it still holds after grading by F . There is a commutative diagram of graded algebras

$$\begin{array}{ccc} & \text{T}_{\mathbb{C}} \text{Der}(\log Y) & \\ \Sigma \swarrow & & \searrow \text{gr } \gamma_Y \\ \text{Sym}_{\mathcal{O}} \text{Der}(\log Y) & \xrightarrow{\alpha_Y} & \text{gr}^F \mathcal{O}[\text{Der}(\log Y)] \\ \pi_Y \downarrow & & \uparrow \\ \text{Rees}_{\mathcal{O}} \text{Der}(\log Y) & \xrightarrow{\cong} & \mathcal{O}[\sigma(\text{Der}(\log Y))] \end{array}$$

Lemma 11. *If α_Y is an isomorphism then*

$$\text{Sym}_{\mathcal{O}} \text{Der}(\log Y) = (i_Y)_* i_Y^{-1} \text{Sym}_{\mathcal{O}} \text{Der}(\log Y)$$

implies $\mathcal{O}[\text{Der}(\log Y)] = (i_Y)_ i_Y^{-1} \mathcal{O}[\text{Der}(\log Y)]$.*

Proof. There is a commutative diagram

$$\begin{array}{ccc}
\mathrm{Sym}_{\mathcal{O}} \mathrm{Der}(\log Y) & \longrightarrow & i_* i^{-1} \mathrm{Sym}_{\mathcal{O}} \mathrm{Der}(\log Y) \\
\downarrow \alpha_Y & & \downarrow i_* i^{-1} \alpha_Y \\
\mathrm{gr}^F \mathcal{O}[\mathrm{Der}(\log Y)] & \xrightarrow{\quad} & i_* i^{-1} \mathrm{gr}^F \mathcal{O}[\mathrm{Der}(\log Y)] \\
& \searrow & \swarrow \\
& \mathrm{gr}^F i_* i^{-1} \mathcal{O}[\mathrm{Der}(\log Y)] &
\end{array}$$

Then the claim follows by induction on $\deg(P)$ for $P \in i_* i^{-1} \mathcal{O}[\mathrm{Der}(\log Y)]$. \square

Lemma 12. α_Y is an isomorphism if and only if π_Y is injective.

Proof. Assume that π_Y is injective. An element of $\mathrm{gr}^F \mathcal{O}[\mathrm{Der}(\log Y)]$ is of the form $\sigma(\gamma_Y(P))$ where $P \in \mathrm{T}_{\mathbb{C}} \mathrm{Der}(\log Y)$. Write $P = P_0 \oplus \cdots \oplus P_d$ where $d = \deg(P)$. If $\sigma(\gamma_Y(P)) \notin \mathrm{im} \alpha_Y$ then $(\mathrm{gr} \gamma_Y)(P_d) = (\mathrm{gr} \gamma_Y)(P) = 0$ and hence $P_d \in \ker \Sigma$ by injectivity of π_Y . By definition of $\mathrm{Sym}_{\mathcal{O}}$, this implies that P_d is in the two-sided ideal generated by the relations $\xi \otimes \eta - \eta \otimes \xi$ and $\xi \otimes (a\eta) - (a\xi) \otimes \eta$ where $\xi, \eta \in \mathrm{Der}(\log Y)$ and $a \in \mathcal{O}$. But

$$\gamma_Y(\xi \otimes \eta - \eta \otimes \xi) = \xi\eta - \eta\xi = [\xi, \eta] \in \mathrm{Der}(\log Y)$$

by involutivity of $\mathrm{Der}(\log Y)$ and

$$\gamma_Y(\xi \otimes (a\eta) - (a\xi) \otimes \eta) = \xi a\eta - a\xi\eta = [\xi, a]\eta = \xi(a)\eta \in \mathrm{Der}(\log Y).$$

This means that

$$\begin{aligned}
\xi \otimes \eta - \eta \otimes \xi &\equiv [\xi, \eta] \pmod{\ker \gamma_Y}, \\
\deg(\xi \otimes \eta - \eta \otimes \xi) &> \deg([\xi, \eta]), \\
\xi \otimes (a\eta) - (a\xi) \otimes \eta &\equiv \xi(a)\eta \pmod{\ker \gamma_Y}, \\
\deg(\xi \otimes (a\eta) - (a\xi) \otimes \eta) &> \deg(\xi(a)\eta).
\end{aligned}$$

Hence $\gamma_Y(P) = \gamma_Y(P')$ and $\deg(P) < \deg(P')$ for some $P' \in \mathrm{T}_{\mathbb{C}} \mathrm{Der}(\log Y)$. Then the claim follows by induction on $d = \deg(P)$. \square

Example 13. Let D and Q be as in Example 9. Then a SINGULAR [GPS05] computation shows that π_D is injective and that

$$\sigma(Q) \notin \alpha_D(\mathrm{Sym}_{\mathcal{O}}^2 \mathrm{Der}(\log D)).$$

By Lemma 12, this implies $Q \notin F_2 \mathcal{O}[\mathrm{Der}(\log D)]$ and hence, by Example 9, $\mathcal{O}[\mathrm{Der}(\log D)] \subsetneq \mathcal{V}_0^D$.

By the following general statement, injectivity of π_Y is equivalent to \mathcal{O} -torsion freeness of $\mathrm{Sym}_{\mathcal{O}} \mathrm{Der}(\log Y)$.

Lemma 14. Let R be a domain and M a finitely presented torsion free R -module. Then the following are equivalent:

- (1) $\mathrm{Sym}_R M$ is R -torsion free.
- (2) $\mathrm{Sym}_R M$ is a domain.
- (3) $\mathrm{Sym}_R M \xrightarrow{\pi_M} \mathrm{Rees}_R M$ is injective.

Proof. Assume that $\mathrm{Sym}_R M$ is R -torsion free. Let $K = Q(R)$ be the fraction field of R . Then $M \otimes_R K \cong K^d$ where $d = \mathrm{rk}(M)$. By choosing a basis of K^d and clearing denominators, one can embed $M \subset R^d$. Then

$$\mathrm{Sym}_R(M) \otimes_R K \cong \mathrm{Sym}_{R \otimes_R K}(M \otimes_R K) \cong \mathrm{Sym}_K(K^d)$$

is a domain and hence $\mathrm{Sym}_R M$ is a domain since R is a domain. Applying Sym_R to the inclusion $M \subset R^d$ yields

$$\begin{array}{ccc} \mathrm{Sym}_R M & \xrightarrow{\phi} & \mathrm{Sym}_R(R^d) \\ & \searrow \pi_M & \nearrow \\ & \mathrm{Rees}_R M & \end{array}$$

and $\ker(\phi) \otimes_R K = 0$ since

$$\mathrm{Sym}_R(M) \otimes_R K \cong \mathrm{Sym}_K(K^d) \cong \mathrm{Sym}_R(R^d) \otimes_R K.$$

A presentation

$$R^m \xrightarrow{(a_{i,j})} R^n \longrightarrow M \longrightarrow 0$$

of M defines an isomorphism

$$\mathrm{Sym}_R M \cong R[T_1, \dots, T_n]/J$$

where $J = \langle \sum_j a_{i,j} T_j \rangle$ is a prime ideal since $\mathrm{Sym}_R M$ is a domain. Since $\mathrm{Sym}_R(R^d)$ is a domain, $\ker \phi$ lifts to a prime ideal $Q \subset R[T_1, \dots, T_n]$. Then $J \subset Q$, $Q \cap R = 0$, and $J \otimes_R K = Q \otimes_R K$ implies $J = Q$ and hence $\ker \pi_M = \ker \phi = 0$. \square

Example 15. Let $D_4 \subset \mathbb{C}^4$ be the central generic hyperplane arrangement defined in Section 5. Then one can compute that the coordinates are zero divisors on $\mathrm{Sym}_{\mathcal{O}}^2 A_4$ at 0. By Lemmata 12, 14, and 24, this implies that α_{D_4} is not an isomorphism.

A divisor $D \subset X$ is called Euler homogeneous if locally $\chi(f) = f$ for some $\chi \in \mathrm{Der}(\log D)$ and $f \in \mathcal{O}$ such that $D = (f)$. In this case, χ is called an Euler vector field and

$$\mathrm{Der}(\log D) \cong \mathcal{O} \cdot \chi \oplus \mathrm{Ann}_{\mathcal{O}}(f).$$

If $\mathrm{Der}(\log D) \cong \mathcal{O} \cdot \chi \oplus A$ then $\mathrm{Sym}_{\mathcal{O}} \mathrm{Der}(\log D) \cong \mathrm{Sym}_{\mathcal{O}}(A)[\chi]$. For an Euler homogeneous divisor D , this implies

$$\mathrm{Sym}_{\mathcal{O}} \mathrm{Der}(\log D) \cong \mathrm{Sym}_{\mathcal{O}}(\mathrm{Ann}_{\mathcal{O}}(f))[\chi].$$

Proposition 16. *Let $D \subset X$ be a divisor such that $\mathrm{Sym}_{\mathcal{O}} \mathrm{Der}(\log D)$ is \mathcal{O} -torsion free. Then $\mathcal{V}_0^D = \mathcal{O}[\mathrm{Der}(\log D)]$ follows from*

$$\mathrm{Sym}_{\mathcal{O}} \mathrm{Der}(\log D) = (i_D)_* i_D^{-1} \mathrm{Sym}_{\mathcal{O}} \mathrm{Der}(\log D).$$

If D is Euler homogeneous and $A = \mathrm{Ann}_{\mathcal{O}}(f)$ or $A \oplus \mathcal{O} \cdot \chi \cong \mathrm{Der}(\log D)$ then the latter is equivalent to $\mathrm{Sym}_{\mathcal{O}} A = (i_D)_ i_D^{-1} \mathrm{Sym}_{\mathcal{O}} A$.*

Proof. This follows from Corollary 5, Lemmata 11, 12, and 14, and the preceding remarks. \square

4. DEPTH AND TORSION OF SYMMETRIC ALGEBRAS

Using a theorem of G. Scheja [Sch61] on extension of coherent analytic sheaves, we shall give sufficient conditions for $\mathcal{V}_0^D = \mathcal{O}[\text{Der}(\log D)]$ in terms of the depth and torsion of the symmetric algebras in Proposition 16.

Theorem 17. *Let $D \subset X$ be a divisor such that $\text{Sym}_{\mathcal{O}} \text{Der}(\log D)$ is \mathcal{O} -torsion free. Let $Z \subset \text{Sing}(D)$ be a closed subset such that $\mathcal{V}_0^D = \mathcal{O}[\text{Der}(\log D)]$ on $X \setminus Z$. Then $\mathcal{V}_0^D = \mathcal{O}[\text{Der}(\log D)]$ on X if*

$$\text{depth}_z(\text{Sym}_{\mathcal{O}}^k \text{Der}(\log D)) \geq \dim_z(Z) + 2$$

for all $z \in Z$ and $k \in \mathbb{N}$. In particular, this holds if D is Euler homogeneous, $A = \text{Ann}_{\Theta}(f)$ or $A \oplus \mathcal{O} \cdot \chi \cong \text{Der}(\log D)$, and

$$\text{depth}_z(\text{Sym}_{\mathcal{O}}^k A) \geq \dim_z(Z) + 2$$

for all $z \in Z$ and $k \in \mathbb{N}$.

Proof. This follows from [Sch61, Satz I-III] and Proposition 16. \square

We shall apply a criterion by C. Huneke [Hun81] for the torsion freeness of symmetric algebras.

Proposition 18. *Let R be a Noetherian domain and let*

$$0 \longrightarrow R \xrightarrow{(a_1, \dots, a_m)^t} R^m \longrightarrow M \longrightarrow 0$$

be a resolution of M . If $\text{grade}(I) \geq k + 1$ for $I = \langle a_1, \dots, a_m \rangle$ then

$$\text{depth}(I, \text{Sym}_R(M)) \geq k.$$

Proof. We proceed by induction on k . By [Hun81, Prop. 2.1], $\text{grade}(I) \geq 2$ implies that $\text{Sym}_R(M)$ is R -torsion free. If $k \geq 2$ then $\text{grade}(I/a) \geq k$ for some $a \in I$ and hence

$$0 \longrightarrow R/a \xrightarrow{([a_1], \dots, [a_m])^t} (R/a)^m \longrightarrow M/a \longrightarrow 0$$

is a resolution of M . Since $\text{Sym}_{R/a}(M/a) \cong \text{Sym}_R(M)/a$, the induction hypothesis applies. \square

Theorem 19. *Let $D \subset X$ be an Euler homogeneous divisor and $A = \text{Ann}_{\Theta}(f)$ or $A \oplus \mathcal{O} \cdot \chi \cong \text{Der}(\log D)$. Let $Z \subset \text{Sing}(D)$ be a closed subset such that $\mathcal{V}_0^D = \mathcal{O}[\text{Der}(\log D)]$ on $X \setminus Z$. For $z \in Z$, let*

$$0 \longrightarrow \mathcal{O}_z \xrightarrow{(a_{z,1}, \dots, a_{z,m})^t} \mathcal{O}_z^m \longrightarrow A_z \longrightarrow 0$$

be a resolution of A_z such that

$$\text{grade}(\langle a_{z,1}, \dots, a_{z,m} \rangle) \geq \dim_z(Z) + 3.$$

Then $\mathcal{V}_0^D = \mathcal{O}[\text{Der}(\log D)]$ on X .

Proof. This follows from Theorem 17, Proposition 18, and [Hun81, Prop. 2.1]. \square

Corollary 20. *Let X be a complex manifold of dimension 3 and let $D \subset X$ be a divisor with only isolated quasi-homogeneous singularities. Then $\mathcal{V}_0^D = \mathcal{O}[\text{Der}(\log D)]$.*

Proof. We may assume that $X \subset \mathbb{C}^3$ is an open neighbourhood of 0, $D = (f)$ with $f \in \mathcal{O}$, and $\text{Sing}(D) = \{0\}$. Let x_1, x_2, x_3 be coordinates on X . Then $\partial(f) = \partial_1(f), \partial_2(f), \partial_3(f) \in \mathfrak{m}_0$ is an \mathcal{O}_0 -sequence. Hence the Koszul-complex

$$\begin{array}{ccccccc}
 0 \rightarrow \mathcal{O}_0 & \xrightarrow{\partial(f)^t} & \mathcal{O}_0^3 & \longrightarrow & \mathcal{O}_0^3 & \longrightarrow & \mathcal{O}_0 \longrightarrow \mathcal{O}_0/\langle \partial(f) \rangle \rightarrow 0 \\
 & & \searrow & & \nearrow & & \\
 & & \text{Ann}_{\mathcal{O}_0}(f) & & & & \\
 & \nearrow & & & \searrow & & \\
 0 & & & & & & 0
 \end{array}$$

is exact and induces a resolution of $\text{Ann}_{\mathcal{O}_0}(f)$. Then the claim follows from Theorem 19 with $Z = \text{Sing}(D)$. \square

Our criterion also applies to some cases of non-isolated singularities.

Example 21. Let $D_3 \subset \mathbb{C}^3$ be the central generic hyperplane arrangement defined in Section 5. Then D_3 is not a free divisor and has a non-isolated singularity at 0. By Lemma 23 and Proposition 24, $A_3 \cong \mathcal{O}^3/\mathcal{O}\cdot(x_1, x_2, x_3)$ and $A_3 \oplus \mathcal{O}\cdot\chi \cong \text{Der}(\log D_3)$. Then, by Examples 1 (2) and 4 (2) on $\text{Sing}(D_3) \setminus \{0\}$ and Theorem 19 for $A = A_3$ and $Z = \{0\}$, $\mathcal{V}_0^{D_3} = \mathcal{O}[\text{Der}(\log D_3)]$.

Our approach may fail in dimension $n > 3$ even for quasi-homogeneous isolated singularities.

Example 22. Let x_1, x_2, x_3, x_4 be coordinates on \mathbb{C}^4 and

$$f = x_1^2 + x_2^2 + x_3^2 + x_4^2.$$

Then $D = (f) \subset \mathbb{C}^4$ has a quasi-homogeneous isolated singularity at 0. One can compute that the coordinates are zero divisors on $\text{Sym}_{\mathcal{O}}^2 \text{Ann}_{\mathcal{O}}(f)$ at 0. By Lemmata 12 and 14 this implies that α_D is not an isomorphism.

5. EXAMPLE OF GENERIC HYPERPLANE ARRANGEMENTS

We shall provide some background for the examples in the previous sections. Let x_1, \dots, x_n be coordinates on \mathbb{C}^n and

$$f_n = x_1 \cdots x_n (x_1 + \cdots + x_n).$$

Then $D_n = (f_n) \subset \mathbb{C}^n$ is a central generic hyperplane arrangement. Let $\chi = \sum_i x_i \partial_i$ be the Euler vector field,

$$\eta_{i,j} = x_i x_j (\partial_i - \partial_j) \in \text{Der}(\log D_n)$$

for $i < j$, and $A_n = \mathcal{O}\langle \eta_{i,j} \rangle$. By J. Wiens [Wie01, Thm. 3.4],

$$\text{Der}(\log D_n) = \mathcal{O}\cdot\chi + A_n$$

with a minimal number of generators. Let

$$\sigma_{i,j,k} = x_i \eta_{j,k} - x_j \eta_{i,k} + x_k \eta_{i,j} \in \text{syz}(\eta_{i,j})$$

for $i < j < k$ and choose a monomial ordering refining $\partial_1 < \cdots < \partial_n$.

Lemma 23. $(\eta_{i,j})$ is a standard basis of A_n and $\text{syz}(\eta_{i,j}) = \langle \sigma_{i,j,k} \rangle$.

Proof. This follows from Buchberger's criterion [GP02, Thm. 1.7.3]. \square

Proposition 24. $\text{Der}(\log D_n) = \mathcal{O}\cdot\chi \oplus A_n$.

Proof. It suffices to verify that no syzygy of χ and the $\eta_{i,j}$ involves χ . One can obtain the syzygies from a standard basis computation [GP02, Alg. 2.5.4]. The first s -polynomials $x_k\chi - \eta_{k,n}$ and $x_j\eta_{i,k} - x_i\eta_{j,k}$ have a zero ∂_n component. Hence only a sequence of s -polynomials starting with $x_k\chi - \eta_{k,n}$ can contribute to syzygies involving χ and the coefficient of χ remains a monomial. Each element in such a sequence has exactly one monomial involving x_n . Since the $\partial_2, \dots, \partial_{n-1}$ are leading components of the $\eta_{i,j}$, the sequence terminates with a non-zero element $a_k\partial_1 \equiv x^{\alpha_k}\chi \pmod{A_n}$. By the same reason, $\mathcal{O} \cdot \partial_1 \oplus A_n$ is a direct sum and hence $x^{\alpha_j}a_k = x^{\alpha_k}a_j$. This implies that the coefficient of χ in any syzygy is zero. \square

REFERENCES

- [CM99] Francisco J. Calderón-Moreno, *Logarithmic differential operators and logarithmic de Rham complexes relative to a free divisor*, Ann. Sci. École Norm. Sup. (4) **32** (1999), no. 5, 701–714. MR MR1710757 (2000g:32010)
- [CMNM05] Francisco Javier Calderón Moreno and Luis Narváez Macarro, *Dualité et comparaison sur les complexes de de Rham logarithmiques par rapport aux diviseurs libres*, Ann. Inst. Fourier (Grenoble) **55** (2005), no. 1, 47–75. MR MR2141288 (2006d:32008)
- [GP02] Gert-Martin Greuel and Gerhard Pfister, *A singular introduction to commutative algebra*, Springer-Verlag, Berlin, 2002, With contributions by Olaf Bachmann, Christoph Lossen and Hans Schönemann, With 1 CD-ROM (Windows, Macintosh, and UNIX). MR MR1930604 (2003k:13001)
- [GPS05] G.-M. Greuel, G. Pfister, and H. Schönemann, *SINGULAR 3.0*, A Computer Algebra System for Polynomial Computations, Centre for Computer Algebra, University of Kaiserslautern, 2005, <http://www.singular.uni-kl.de>.
- [Hun81] Craig Huneke, *On the symmetric algebra of a module*, J. Algebra **69** (1981), no. 1, 113–119. MR MR613861 (82d:13016)
- [Meb89] Z. Mebkhout, *Le formalisme des six opérations de Grothendieck pour les D_X -modules cohérents*, Travaux en Cours [Works in Progress], vol. 35, Hermann, Paris, 1989, With supplementary material by the author and L. Narváez Macarro. MR MR1008245 (90m:32026)
- [Sai80] Kyoji Saito, *Theory of logarithmic differential forms and logarithmic vector fields*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **27** (1980), no. 2, 265–291. MR MR586450 (83h:32023)
- [Sch61] Günter Scheja, *Riemannsche Hebbbarkeitssätze für Cohomologieklassen*, Math. Ann. **144** (1961), 345–360. MR MR0148941 (26 #6437)
- [Tor04] Tristan Torrelli, *On meromorphic functions defined by a differential system of order 1*, Bull. Soc. Math. France **132** (2004), no. 4, 591–612. MR MR2131905 (2005m:32015)
- [Wie01] Jonathan Wiens, *The module of derivations for an arrangement of subspaces*, Pacific J. Math. **198** (2001), no. 2, 501–512. MR MR1835521 (2002d:14090)

MATHIAS SCHULZE, DEPARTMENT OF MATHEMATICS, OKLAHOMA STATE UNIVERSITY, 401 MSCS, STILLWATER, OK 74078, USA

E-mail address: mschulze@math.okstate.edu