

MATROID CONNECTIVITY AND SINGULARITIES OF CONFIGURATION HYPERSURFACES

GRAHAM DENHAM, MATHIAS SCHULZE, AND ULI WALTHER

ABSTRACT. Consider a linear realization of a matroid over a field. One associates with it a configuration polynomial and a symmetric bilinear form with linear homogeneous coefficients. The corresponding configuration hypersurface and its non-smooth locus support the respective first and second degeneracy scheme of the bilinear form.

We show that these schemes are reduced and describe the effect of matroid connectivity: for (2-)connected matroids, the configuration hypersurface is integral, and the second degeneracy scheme is reduced Cohen–Macaulay of codimension 3. If the matroid is 3-connected, then also the second degeneracy scheme is integral.

In the process, we describe the behavior of configuration polynomials, forms and schemes with respect to various matroid constructions.

CONTENTS

1. Introduction	2
1.1. Feynman diagrams	2
1.2. Configuration polynomials	3
1.3. Summary of results	4
1.4. Outline of the proof	5
Acknowledgments	7
2. Matroids and configurations	8
2.1. Matroid basics	8
2.2. Handle decomposition	11
2.3. Configurations and realizations	15
2.4. Graphic matroids	18
3. Configuration polynomials and forms	22
3.1. Configuration polynomials	22

2010 *Mathematics Subject Classification*. Primary 14N20; Secondary 05C31, 14B05, 14M12, 81Q30.

Key words and phrases. Configuration, matroid, singularity, Feynman, Kirchhoff, Symanzik, Cohen–Macaulay, determinantal.

GD was supported by NSERC of Canada. MS was supported by the German Research Foundation (DFG) in the collaborative research center TRR 195 under Project B05 (#324841351). UW was supported by the NSF Grant DMS-1401392 and by the Simons Foundation Collaboration Grant for Mathematicians #580839.

3.2. Graph polynomials	32
3.3. Configuration forms	33
4. Configuration hypersurfaces	38
4.1. Commutative ring basics	38
4.2. Jacobian and degeneracy schemes	42
4.3. Generic points and codimension	46
4.4. Reducedness of degeneracy schemes	49
4.5. Integrality of degeneracy schemes	55
5. Examples	59
References	61

1. INTRODUCTION

1.1. Feynman diagrams. A basic problem in high-energy physics is to understand the scattering of particles. The basic tool for theoretical predictions is the Feynman diagram with underlying Feynman graph $G = (V, E)$. The scattering data correspond to Feynman integrals, computed in the positive orthant of the projective space labeled by the internal edges of the Feynman graph. The integrand is the square root of a rational function in the edge variables x_e , $e \in E$, that depends parametrically on the masses and moments of the involved particles (see [Bro17]).

The convergence of a Feynman integral is determined by the structure of the denominator of this rational function, which always involves a power of the square root of the *Symanzik polynomial* $\sum_{T \in \mathcal{T}_G} \prod_{e \notin T} x_e$ of G where \mathcal{T}_G denotes the set of spanning trees of G . The remaining factor of the denominator, appearing for graphs with edge number less than twice the loop number, is a power of the square root of the second Symanzik polynomial obtained by summing over 2-forests and involves masses and moments. Symanzik polynomials can factor, and the singularities and intersections of the individual components determine the behavior of the Feynman integrals.

Until about a decade ago, all explicitly computed integrals were built from multiple Riemann zeta values and polylogarithms; for example, Broadhurst and Kreimer display a large body of such computations in [BK97]. In fact, Kontsevich at some point speculated that Symanzik polynomials, or equivalently their cousins the *Kirchhoff polynomials*

$$\psi_G(x) = \sum_{T \in \mathcal{T}_G} \prod_{e \in T} x_e$$

be mixed Tate; this would imply the relation to multiple zeta values. However, Belkale and Brosnan [BB03] proved that the collection of Kirchhoff polynomials is a rather complicated class of singularities: their hypersurface complements generate the ring of all geometric motives. This does not exactly rule out that Feynman integrals are in

some way well-behaved, but makes it rather less likely, and explicit counterexamples to Kontsevich’s conjecture were subsequently worked out by Doryn [Dor11] as well as by Brown and Schnetz [BS12]. On the other hand, these examples make the study of these singularities, and especially any kind of uniformity results, that much more interesting.

The influential paper [BEK06] of Bloch, Esnault and Kreimer generated a significant amount of work from the point of view of complex geometry: we refer to the book [Mar10] of Marcolli for exposition, as well as [Bro17; Dor11; BW10]. Varying ideas of Connes and Kreimer on renormalization that view Feynman integrals as specializations of the Tutte polynomial, Aluffi and Marcolli formulate in [AM11b; AM11a] parametric Feynman integrals as periods, leading to motivic studies on cohomology. On the explicit side, there is a large body of publications in which specific graphs and their polynomials and Feynman integrals are discussed. But, as Brown writes in [Bro15], while a diversity of techniques is used to study Feynman diagrams, “each new loop order involves mathematical objects which are an order of magnitude more complex than the last, [...] the unavoidable fact is that arbitrary integrals remain out of reach as ever.”

The present article can be seen as the first step towards a search for uniform properties in this zoo of singularities. We view it as a stepping stone for further studies of invariants such as log canonical threshold, logarithmic differential forms and embedded resolution of singularities.

1.2. Configuration polynomials. The main idea of Belkale and Brosnan is to move the burden of proof into the more general realm of polynomials and constructible sets derived from matroids rather than graphs, and then to reduce to known facts about such polynomials. The article [BEK06] casts Kirchhoff and Symanzik polynomials as very special instances of *configuration polynomials*; this idea was further developed by Patterson in [Pat10]. We consider this as a more natural setting since notions such as duality and quotients behave well for configuration polynomials as a whole, but these operations do not preserve the subfamily of matroids derived from graphs. In particular, we can focus exclusively on Kirchhoff/configuration polynomials, since the Symanzik polynomial of G appears as the configuration polynomial of the dual configuration induced by the incidence matrix of G .

The configuration polynomial does not depend on a matroid itself but on a configuration, that is, on a (linear) realization of a matroid over a field \mathbb{K} . The same matroid can admit different realizations, which, in turn, give rise to different configuration polynomials (see Example 5.3). The *matroid (basis) polynomial* is a competing object, which is assigned to any, even non-realizable, matroid. It has proven useful for combinatorial applications (see [AOV18; Piq19]). For graphs and, more generally, regular matroids, all configuration polynomials

essentially agree with the matroid polynomial. In general, however, configuration polynomials differ significantly from matroid polynomials, as documented in Example 5.2.

Configuration polynomials have a geometric feature that matroid polynomials lack: generalizing Kirchhoff's matrix-tree theorem, the configuration polynomial arises as the determinant of a symmetric bilinear *configuration form* with linear polynomial coefficients. As a consequence, the corresponding *configuration hypersurface* maps naturally to the generic symmetric determinantal variety. In the present article, we establish further uniform, geometric properties of configuration polynomials, which we observe do not hold for matroid polynomials in general.

1.3. Summary of results. Some indication of what is to come can be gleaned from the following note by Marcolli in [Mar10, p. 71]: “graph hypersurfaces tend to have singularity loci of small codimension”.

Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid \mathbf{M} of rank $\text{rk } \mathbf{M} = \dim W$ on a set E (see Definition 2.14). Fix coordinates $x_E = (x_e)_{e \in E}$. There is an associated symmetric *configuration (bilinear) form* Q_W with linear homogeneous coefficients (see Definition 3.20). Its determinant is the *configuration polynomial* (see Definition 3.2 and Lemma 3.23)

$$\psi_W = \det Q_W = \sum_{B \in \mathcal{B}_M} c_{W,B} \cdot \prod_{e \in B} x_e \in \mathbb{K}[x_E]$$

where \mathcal{B}_M denotes the set of bases of \mathbf{M} and the coefficients $c_{W,B} \in \mathbb{K}^*$ depend of the realization W . The *configuration hypersurface* defined by ψ_W is the scheme

$$X_W = \text{Spec}(\mathbb{K}[x_E]/\langle \psi_W \rangle) \subseteq \mathbb{K}^E.$$

It can be seen as the *first degeneracy scheme* of Q_W (see Definition 4.9). The *second degeneracy scheme* $\Delta_W \subseteq \mathbb{K}^E$ of Q_W , defined by the submaximal minors of Q_W , is a subscheme of the *Jacobian scheme* $\Sigma_W \subseteq \mathbb{K}^E$ of X_W , defined by ψ_W and its partial derivatives (see Lemma 4.12). The latter defines the non-smooth locus of X_W over \mathbb{K} , which is the singular locus of X_W if \mathbb{K} is perfect (see Remark 4.10). Paterson showed Σ_W and Δ_W have the same underlying reduced scheme (see Theorem 4.17), that is,

$$\Delta_W \subseteq \Sigma_W \subseteq \mathbb{K}^E, \quad \Sigma_W^{\text{red}} = \Delta_W^{\text{red}}.$$

We give a simple proof of this fact. He mentions that he does not know the reduced scheme structure (see [Pat10, p. 696]). We show that Σ_W is typically not reduced (see Example 5.1), whereas Δ_W often is. Our main results from Theorems 4.16, 4.25, 4.36 and 4.37 can be summarized as follows.

Main Theorem. *Let \mathbf{M} be a matroid on the set E with a linear realization $W \subseteq \mathbb{K}^E$ over a field \mathbb{K} . Then the configuration hypersurface*

X_W is reduced and generically smooth over \mathbb{K} . Moreover, the second degeneracy scheme Δ_W is also reduced and agrees with Σ_W^{red} , the non-smooth locus of X_W over \mathbb{K} . Unless \mathbb{K} has characteristic 2, the Jacobian scheme Σ_W is generically reduced.

Suppose now that \mathbf{M} is connected. Then X_W is integral unless \mathbf{M} has rank zero. Suppose in addition that the rank of \mathbf{M} is at least 2. Then Δ_W is Cohen–Macaulay of codimension 3 in \mathbb{K}^E . If, moreover, \mathbf{M} is 3-connected, then Δ_W is integral. \square

Note that $X_W = \emptyset$ if $\text{rk } \mathbf{M} = 0$ and $\Sigma_W = \emptyset = \Delta_W$ if $\text{rk } \mathbf{M} \leq 1$ (see Remarks 3.5 and 4.13.(a)). It suffices to require the connectedness hypotheses after deleting all loops (see Remark 4.11). If \mathbf{M} is disconnected even after deleting all loops, then Σ_W and hence Δ_W has codimension 2 in \mathbb{K}^E (see Proposition 4.16).

While our main objective is to establish the results above, along the way we continue the systematic study of configuration polynomials in the spirit of [BEK06; Pat10]. For instance, we describe the behavior of configuration polynomials with respect to connectedness, duality, deletion/contraction and 2-separations (see Propositions 3.8, 3.10, 3.12 and 3.27). Patterson showed that the *second Symanzik polynomial* associated with a Feynman graph is, in fact, a configuration polynomial. More precisely, we explain that its dual, the *second Kirchhoff polynomial*, is associated with the quotient of the graph configuration by the momentum parameters (see Proposition 3.19). In this way, Patterson’s result becomes a special case of a formula for configuration polynomials of elementary quotients (see Proposition 3.14).

1.4. Outline of the proof. The proof of the Main Theorem intertwines methods from matroid theory, commutative algebra and algebraic geometry. In order to keep our arguments self-contained and accessible, we recall preliminaries from each of these subjects and give detailed proofs (see §2.1, §2.3 and §4.1). One easily reduces the claims to the case where \mathbf{M} is connected (see Proposition 3.8 and Theorem 4.36).

An important commutative algebra ingredient is a result of Kutz (see [Kut74]): the grade of an ideal of submaximal minors of a symmetric matrix cannot exceed 3, and equality forces the ideal to be perfect. Kutz’ result applies to the defining ideal of Δ_W . The codimension of Δ_W in \mathbb{K}^E is therefore bounded by 3 and Δ_W is Cohen–Macaulay in case of equality (see Proposition 4.19). In this case, Δ_W is pure-dimensional, and hence, it is reduced if it is generically reduced (see Lemma 4.4).

On the matroid side our approach makes use of *handles* (see Definition 2.3), which are called *ears* in case of graphic matroids. A *handle decomposition* builds up any connected matroid from a circuit by successively attaching handles (see Definition 2.6). Conversely, this yields for any connected matroid which is not a circuit a *non-disconnective*

handle which leaves the matroid connected when deleted (see Definition 2.3). This allows one to prove statements on connected matroids by induction.

We describe the effect of deletion and contraction of a handle H to the configuration polynomial (see Corollary 3.13). In case the Jacobian scheme $\Sigma_{W \setminus H}$ associated with the deletion $\mathbf{M} \setminus H$ has codimension 3 we prove the same for Σ_W (see Lemma 4.22). Applied to a non-disconnective H it follows with Patterson's result that Δ_W reaches the dimension bound and is thus Cohen–Macaulay of codimension 3 (see Theorem 4.25). We further identify three (more or less explicit) types of generic points with respect to a non-disconnective handle (see Corollary 4.26).

In case $\text{ch } \mathbb{K} \neq 2$, generic reducedness of Σ_W implies (generic) reducedness of Δ_W . The schemes Σ_W and Δ_W show similar behavior with respect to deletion and contraction (see Lemmas 4.29 and 4.31). As a consequence, generic reducedness can be proved along the same lines (see Lemma 4.35). In both cases, we have to show reducedness at all (the same) generic points. In what follows, we restrict ourselves to Δ_W . Our proof proceeds by induction on the cardinality $|E|$ of the underlying set E of the matroid \mathbf{M} .

Unless \mathbf{M} a circuit, the handle decomposition guarantees the existence of a non-disconnective handle H . In case $H = \{h\}$ has size 1, the scheme $\Delta_{W \setminus h}$ associated with the deletion $\mathbf{M} \setminus h$ is the intersection of Δ_W with the divisor x_e (see Lemma 4.29). This serves to recover generic reducedness of Δ_W from $\Delta_{W \setminus h}$ (see Lemma 4.30). The same argument works if H does not arise from a handle decomposition.

This leads us to consider non-disconnective handles independently of a handle decomposition. They turn out to be special instances of maximal handles which form the *handle partition* of the matroid (see Lemma 2.4). As a purely matroid-theoretic ingredient, we show that the number of non-disconnective handles is strictly increasing when adding handles (see Proposition 2.12). For handle decompositions of length 2, a distinguished role is played by the prism matroid (see Example 2.7). Its handle partition consists of 3 non-disconnective handles of size 2 (see Lemmas 2.10 and 2.25). Here an explicit calculation shows that Δ_W is reduced in the torus $(\mathbb{K}^*)^6$ (see Lemma 4.28). The corresponding result for Σ_W holds only if $\text{ch } \mathbb{K} \neq 2$.

Suppose now that \mathbf{M} is not a circuit and has no non-disconnective handles of size 1. Then $\mathbf{M} \setminus e$ might be disconnected for all $e \in E$ and does not qualify for an inductive step. In this case, we aim instead for contracting W by a suitable subset $G \subsetneq E$ which keeps \mathbf{M} connected. In the partial torus $\mathbb{K}^F \times (\mathbb{K}^*)^G$ where $F := E \setminus G$, the scheme $\Delta_{W/G}$ associated with the contraction \mathbf{M}/G relates to the *normal cone* of Δ_W along the coordinate subspace $V(x_F)$ where $x_F = (x_f)_{f \in F}$ (see Lemma 4.31). To induce generic reducedness from $\Delta_{W/G}$ to Δ_W , we

pass through a *deformation to the normal cone*, which is our main ingredient from algebraic geometry. The role of x_h above is then played by the deformation parameter t .

In algebraic terms, this deformation is represented by the *Rees algebra* $\text{Rees}_I R$ with respect to an ideal $I \trianglelefteq R$, and the normal cone by the *associated graded ring* $\text{gr}_I R$ (see Definition 4.6). Passing through $\text{Rees}_I R$, we recover generic reducedness of R along $V(I)$ from generic reducedness of $\text{gr}_I R$ (see Definition 4.3 and Lemma 4.7). By assumption on \mathbf{M} , there are at least 3 more elements in E than maximal handles (see Proposition 2.12), and \mathbf{M} is the prism matroid in case of equality. Based on a strict inequality, we use a codimension argument to construct a suitable partition $E = F \sqcup G$ for which *all* generic points of Δ_W are *along* $V(x_F)$ (see Lemma 4.34). This yields generic reducedness of Δ_W in this case (see Lemma 4.32). A slight modification of the approach also covers the generic points outside the torus $(\mathbb{K}^*)^6$ if \mathbf{M} is the prism matroid. The case where \mathbf{M} is a circuit is reduced to that where \mathbf{M} is a triangle by successively contracting an element of E (see Lemma 4.33). In this base case Δ_W is a reduced point, but Σ_W is reduced only if $\text{ch } \mathbb{K} \neq 2$ (see Example 4.14).

Finally, suppose that \mathbf{M} is a 3-connected matroid. Here we prove that Δ_W is irreducible and hence integral, which implies that Σ is irreducible (see Theorem 4.37). We first observe that handles of (co)size at least 2 are 2-separations (see Lemma 2.4.(e)). It follows that the handle decomposition consists entirely of non-disconnective 1-handles (see Proposition 2.5) and that all generic points of Δ_W lie in the torus $(\mathbb{K}^*)^E$ (see Corollary 4.27). We show that the number of generic points is bounded by that of $\Delta_{W \setminus e}$ for all $e \in E$ (see Lemma 4.30). Duality switches deletion and contraction and identifies generic points of Δ_W and Δ_{W^\perp} (see Corollary 4.18). Using Tutte's wheels-and-whirls theorem, the irreducibility of Δ_W can therefore be reduced to the cases where \mathbf{M} is a wheel W_n or a whirl W^n for some $n \geq 3$ (see Example 2.26 and Lemma 4.38). For fixed n , we show that the schemes X_W , Σ_W and Δ_W are all isomorphic for all realizations W of W_n and W^n (see Proposition 4.40). An induction on n with an explicit study of the base cases $n \leq 4$ finishes the proof (see Corollary 4.41 and Lemma 4.43).

Acknowledgments. The project whose results are presented here started with a research in pairs meeting at the Centro de Giorgi in Pisa in February 2018. We thank the Institute for a pleasant stay in a stimulating research environment. We also thank Aldo Conca, Raul Epure, Darij Grinberg, Delphine Pol and Karen Yeats for helpful comments. We are grateful to the referees for a careful reading of the manuscript and resulting improvements to the exposition.

This is a preprint of an article published in Letters in Mathematical Physics. The final authenticated version is available online at: <https://doi.org/10.1007/s11005-020-01352-3>

2. MATROIDS AND CONFIGURATIONS

Our algebraic objects of interest are associated with a realization of a matroid. In this section, we prepare the path for an inductive approach driven by the underlying matroid structure. Our main tool is the handle decomposition, a matroid version of the ear decomposition of graphs.

2.1. Matroid basics. In this subsection, we review the relevant basics of matroid theory using Oxley's book (see [Oxl11]) as a comprehensive reference.

Denote by $\text{Min } \mathcal{P}$ and $\text{Max } \mathcal{P}$ the set of minima and maxima of a poset \mathcal{P} . Let \mathbf{M} be a *matroid* on a set $E =: E_{\mathbf{M}}$. We use this font throughout to denote matroids. With 2^E partially ordered by inclusion, \mathbf{M} can be defined by a monotone submodular *rank* function (see [Oxl11, Cor. 1.3.4])

$$\text{rk} = \text{rk}_{\mathbf{M}}: 2^E \rightarrow \mathbb{N} = \{0, 1, 2, \dots\}$$

with $\text{rk}(S) \leq |S|$ for any subset $S \subseteq E$. The *rank* of \mathbf{M} is then

$$\text{rk } \mathbf{M} := \text{rk}_{\mathbf{M}}(E).$$

Alternatively, it can be defined in terms of each of the following collections of subsets of E (see [Oxl11, Prop. 1.3.5, p. 28]):

- *independent sets* $\mathcal{I}_{\mathbf{M}} = \{I \subseteq E \mid |I| = \text{rk}_{\mathbf{M}}(I)\} \subseteq 2^E$,
- *bases* $\mathcal{B}_{\mathbf{M}} = \text{Max } \mathcal{I}_{\mathbf{M}} = \{B \subseteq E \mid |B| = \text{rk}_{\mathbf{M}}(B) = \text{rk } \mathbf{M}\} \subseteq 2^E$,
- *circuits* $\mathcal{C}_{\mathbf{M}} = \text{Min}(2^E \setminus \mathcal{I}_{\mathbf{M}}) \subseteq 2^E$,
- *flats* $\mathcal{L}_{\mathbf{M}} = \{F \subseteq E \mid \forall e \in E \setminus F: \text{rk}_{\mathbf{M}}(F \cup \{e\}) > \text{rk}_{\mathbf{M}}(F)\}$.

For instance (see [Oxl11, Lem. 1.3.3]), for any subset $S \subseteq E$,

$$(2.1) \quad \text{rk}_{\mathbf{M}}(S) = \max \{|I| \mid S \supseteq I \in \mathcal{I}_{\mathbf{M}}\}.$$

The *closure* operator of \mathbf{M} is defined by (see [Oxl11, Lem. 1.4.2])

$$(2.2) \quad \text{cl}_{\mathbf{M}}: 2^E \mapsto \mathcal{L}_{\mathbf{M}}, \quad \text{rk}_{\mathbf{M}} = \text{rk}_{\mathbf{M}} \circ \text{cl}_{\mathbf{M}}.$$


The following matroid plays a special role in the proof of our main result.

Definition 2.1 (Prism matroid). The *prism matroid* has underlying set E with $|E| = 6$ and circuits

$$\mathcal{C}_{\mathbf{M}} = \{\{e_1, e_2, e_3, e_4\}, \{e_1, e_2, e_5, e_6\}, \{e_3, e_4, e_5, e_6\}\}.$$

The name comes from the observation that its independent sets $\mathcal{I}_{\mathbf{M}}$ are the affinely independent subsets of the vertices of the triangular prism (see Figure 1).

FIGURE 1. The triangular prism.



The elements of $E \setminus \bigcup \mathcal{B}_M$ and $\bigcap \mathcal{B}_M$ are called *loops* and *coloops* in M , respectively. A matroid is *free* if $E \in \mathcal{B}_M$, that is, every $e \in E$ is a coloop in M . By a k -*circuit* in M we mean a circuit $C \in \mathcal{C}_M$ with $|C| = k$ elements, 3-circuits are called *triangles*.

The circuits in M give rise to an equivalence relation on E by declaring $e, f \in E$ equivalent if $e = f$ or $e, f \in C$ for some $C \in \mathcal{C}_M$ (see [Oxl11, Prop. 4.1.2]). The corresponding equivalence classes are the *connected components* of M . If there is at most one such a component, then M is said to be *connected*. The *connectivity function* of M is defined by

$$\lambda_M: 2^E \rightarrow \mathbb{N}, \quad \lambda_M(S) := \text{rk}_M(S) + \text{rk}_M(E \setminus S) - \text{rk}(M).$$

For $k \geq 1$, a subset $S \subseteq E$, or the partition $E = S \sqcup (E \setminus S)$, is called a k -*separation* of M if

$$\lambda_M(S) < k \leq \min \{|S|, |E \setminus S|\}.$$

It is called *exact* if the latter is an equality. The *connectivity* $\lambda(M)$ of M is the minimal k for which there is a k -separation of M , or $\lambda(M) = \infty$ if no such exists. The matroid M is said to be k -*connected* if $\lambda(M) \geq k$. Connectedness is the special case $k = 2$.

We now review some standard constructions of new matroids from old. Their geometric significance is explained in §2.3.

The *direct sum* $M_1 \oplus M_2$ of matroids M_1 and M_2 is the matroid on $E_{M_1} \sqcup E_{M_2}$ with independent sets

$$(2.3) \quad \mathcal{I}_{M_1 \oplus M_2} := \{I_1 \sqcup I_2 \mid I_1 \in \mathcal{I}_{M_1}, I_2 \in \mathcal{I}_{M_2}\}.$$

The sum is *proper* if $E_{M_1} \neq \emptyset \neq E_{M_2}$. Connectedness means that a matroid is not a proper direct sum (see [Oxl11, Prop. 4.2.7]). In particular, any (co)loop is a connected component.

Let $F \subseteq E$ be any subset. Then the *restriction matroid* $M|_F$ is the matroid on F with independent sets and bases (see [Oxl11, 3.1.12, 3.1.14])

$$(2.4) \quad \mathcal{I}_{M|_F} = \mathcal{I}_M \cap 2^F, \quad \mathcal{B}_{M|_F} = \text{Max} \{B \cap F \mid B \in \mathcal{B}_M\}.$$

Its set of circuits is (see [Oxl11, 3.1.13])

$$(2.5) \quad \mathcal{C}_{M|_F} = \mathcal{C}_M \cap 2^F.$$

By definition, $\text{rk}_{M|_F} = \text{rk}_M|_{2^F}$, so we may omit the index without ambiguity. Thinking of restriction to $E \setminus F$ as an operation that deletes elements in F from E , one defines the *deletion matroid*

$$M \setminus F := M|_{E \setminus F}.$$

The *contraction matroid* M/F is the matroid on $E \setminus F$ with independent sets and bases (see [Oxl11, Prop. 3.1.7, Cor. 3.1.8])

$$(2.6) \quad \begin{aligned} \mathcal{I}_{M/F} &= \{I \subseteq E \setminus F \mid I \cup B \in \mathcal{I}_M \text{ for some/every } B \in \mathcal{B}_{M|_F}\}, \\ \mathcal{B}_{M/F} &= \{B' \subseteq E \setminus F \mid B' \cup B \in \mathcal{B}_M \text{ for some/every } B \in \mathcal{B}_{M|_F}\}. \end{aligned}$$

Its circuits are the minimal non-empty sets $C \setminus F$ where $C \in \mathcal{C}_M$ (see [Oxl11, Prop. 3.1.10]), that is,

$$(2.7) \quad \mathcal{C}_{M/F} = \text{Min} \{C \setminus F \mid F \not\subseteq C \in \mathcal{C}_M\}.$$

In §2.3, E will be a basis and E^\vee the corresponding dual basis. We often identify $E = E^\vee$ by the bijection

$$(2.8) \quad \nu: E \rightarrow E^\vee, \quad e \mapsto e^\vee.$$

The complement of a subset $S \subseteq E$ corresponds to

$$S^\perp := \nu(E \setminus S) \subseteq E^\vee.$$

The *dual matroid* M^\perp is the matroid on E^\vee with bases

$$(2.9) \quad \mathcal{B}_{M^\perp} = \{B^\perp \mid B \in \mathcal{B}_M\}.$$

In particular, we have (see [Oxl11, 2.1.8])

$$\text{rk } M + \text{rk } M^\perp = |E|.$$

Connectivity is invariant under dualizing (see [Oxl11, Cor. 8.1.5]),

$$(2.10) \quad \lambda_M = \lambda_{M^\perp} \circ \nu, \quad \lambda(M) = \lambda(M^\perp).$$

We use ν^{-1} in place of (2.8) for M^\perp , so that $S^{\perp\perp} = S$. For subsets $F \subseteq E$ and $G \subseteq E^\vee$, one can identify (see [Oxl11, 3.1.1])

$$(2.11) \quad \begin{aligned} (M/F)^\perp &= M^\perp|_{F^\perp} = M^\perp \setminus \nu(F), \\ (M \setminus \nu^{-1}(G))^\perp &= (M|_{G^\perp})^\perp = M^\perp/G. \end{aligned}$$

Various matroid data of M^\perp is also considered as *codata* of M . A *triad* of M is a 3-cocircuit of M , that is, a triangle of M^\perp .

Example 2.2 (Uniform matroids). The *uniform matroid* $U_{r,n}$ of rank $r \geq 0$ on a set E of size $|E| = n$ has bases

$$\mathcal{B}_{U_{r,n}} = \{B \subseteq E \mid |B| = r\}.$$

For $r = n$ it is the free matroid of rank r . It is connected if and only if $0 < r < n$. By definition, $U_{r,n}^\perp = U_{n-r,n}$ for all $0 \leq r \leq n$.

Informally, we refer to a matroid \mathbf{M} on E for which $E \in \mathcal{C}_{\mathbf{M}}$, and hence, $\mathcal{C}_{\mathbf{M}} = \{E\}$, as a *circuit*, and as a *triangle* if $|E| = 3$. It is easily seen that such a matroid is $\mathbf{U}_{n-1,n}$ where $n = |E|$, and that $\lambda(\mathbf{U}_{n-1,n}) = 2$. \diamond

2.2. Handle decomposition. In this subsection, we investigate handles as building blocks of connected matroids.

Definition 2.3 (Handles). Let \mathbf{M} be a matroid. A subset $\emptyset \neq H \subseteq E$ is a *handle* in \mathbf{M} if $C \cap H \neq \emptyset$ implies $H \subseteq C$ for all $C \in \mathcal{C}_{\mathbf{M}}$. Write $\mathcal{H}_{\mathbf{M}}$ for the set of handles in \mathbf{M} , ordered by inclusion. A *subhandle* of $H \in \mathcal{H}_{\mathbf{M}}$ is a subset $\emptyset \neq H' \subseteq H$. We call $H \in \mathcal{H}_{\mathbf{M}}$

- *proper* if $H \neq E$,
- *maximal* if $H \in \text{Max } \mathcal{H}_{\mathbf{M}}$,
- a *k-handle* if $|H| = k$,
- *disconnective* if $\mathbf{M} \setminus H$ is disconnected and
- *separating* if $\min\{|H|, |E \setminus H|\} \geq 2$.

Singletons $\{e\}$ and subhandles are handles. If $\bigcup \mathcal{C}_{\mathbf{M}} \neq E$, then $E \setminus \bigcup \mathcal{C}_{\mathbf{M}} \in \text{Max } \mathcal{H}_{\mathbf{M}}$ and is a union of coloops. The maximal handles in $\bigcup \mathcal{C}_{\mathbf{M}}$ are the minimal non-empty intersections of all subsets of $\mathcal{C}_{\mathbf{M}}$. Together they form the *handle partition* of E

$$E = \bigsqcup_{H \in \text{Max } \mathcal{H}_{\mathbf{M}}} H,$$

which refines the partition of $\bigcup \mathcal{C}_{\mathbf{M}}$ into connected components. In particular, each circuit is a disjoint union of maximal handles. For any subset $F \subseteq E$, (2.5) yields an inclusion

$$\mathcal{H}_{\mathbf{M}} \cap 2^F \subseteq \mathcal{H}_{\mathbf{M}|_F}.$$

Lemma 2.4 (Handle basics). *Let \mathbf{M} be a matroid and $H \in \mathcal{H}_{\mathbf{M}}$.*

- (a) *If $H = E$, then $\mathbf{M} = \mathbf{U}_{r,n}$ where $n = |E| \geq 1$ and $r \in \{n-1, n\}$ (see Example 2.2). In the latter case, $|E| = 1$ if \mathbf{M} is connected.*
- (b) *Either $H \in \mathcal{I}_{\mathbf{M}}$ or $H \in \mathcal{C}_{\mathbf{M}}$. In the latter case, H is maximal and a connected component of \mathbf{M} . In particular, if \mathbf{M} is connected and H is proper, then $H \in \mathcal{I}_{\mathbf{M}}$, $H \subsetneq C$ for some circuit $C \in \mathcal{C}_{\mathbf{M}}$, and $H \in \mathcal{C}_{\mathbf{M}/(E \setminus H)}$.*
- (c) *For any subhandle $\emptyset \neq H' \subseteq H$, $H \setminus H'$ consists of coloops in $\mathbf{M} \setminus H'$. In particular, non-disconnective handles are maximal.*
- (d) *If $H \notin \mathcal{C}_{\mathbf{M}}$, then there is a bijection*

$$\mathcal{C}_{\mathbf{M}} \rightarrow \mathcal{C}_{\mathbf{M}/H}, \quad C \mapsto C \setminus H.$$

If $H \notin \text{Max } \mathcal{H}_{\mathbf{M}}$, then there is a bijection

$$\text{Max } \mathcal{H}_{\mathbf{M}} \rightarrow \text{Max } \mathcal{H}_{\mathbf{M}/H}, \quad H' \mapsto H' \setminus H,$$

which identifies non-disconnective handles. In this case, the connected components of \mathbf{M} which are not contained in $H \setminus \bigcup \mathcal{C}_{\mathbf{M}}$ correspond to the connected components of \mathbf{M}/H .

(e) Suppose that \mathbf{M} is connected and H is proper. Then

$$\mathrm{rk}(\mathbf{M}/H) = \mathrm{rk} \mathbf{M} - |H|, \quad \lambda_{\mathbf{M}}(H) = 1.$$

In particular, if H is separating, then H is a 2-separation of \mathbf{M} .

Proof.

(a) Suppose that $H = E$. Then $\mathcal{C}_{\mathbf{M}} \subseteq \{E\}$ and $\mathbf{M} = \mathbf{U}_{n-1,n}$ in case of equality. Otherwise, $\mathcal{C}_{\mathbf{M}} = \emptyset$ implies $\mathcal{B}_{\mathbf{M}} = \{E\}$ and $\mathbf{M} = \mathbf{U}_{n,n}$ (see [Oxl11, Prop. 1.1.6]).

(b) Suppose that $H \notin \mathcal{I}_{\mathbf{M}}$. Then there is a circuit $H \supseteq C \in \mathcal{C}_{\mathbf{M}}$. By definition of handle and incomparability of circuits, $H = C \setminus (E \setminus H) \in \mathcal{C}_{\mathbf{M}/(E \setminus H)}$ (see (2.7)) and $H = C$ is disjoint from all other circuits and hence a connected component of \mathbf{M} .

(c) Suppose that $h \in H \setminus H'$ is not a coloop in $\mathbf{M} \setminus H'$. Then $h \in C \cap H$ for some $C \in \mathcal{C}_{\mathbf{M} \setminus H'} \subseteq \mathcal{C}_{\mathbf{M}}$ (see (2.5)) and hence $H' \subseteq H \subseteq C$ since H is a handle, a contradiction.

(d) The first bijection follows from (2.7) with $F = H$. The remaining claims follow from the discussion preceding the lemma.

(e) Part (b) yields the first equality (see [Oxl11, Prop. 3.1.6]) along with a circuit $H \neq C \in \mathcal{C}_{\mathbf{M}}$. Pick a basis $B \in \mathcal{B}_{\mathbf{M} \setminus H}$. Clearly $S := B \sqcup H$ spans \mathbf{M} . For any $h \in H$, we check that $S \setminus \{h\} \in \mathcal{I}_{\mathbf{M}}$. Otherwise, there is a circuit $S \setminus \{h\} \supseteq C \in \mathcal{C}_{\mathbf{M}}$. Since $C \not\subseteq B$ and by definition of handle, we have $H \cap C \neq \emptyset$ and hence $h \in H \subseteq C$, a contradiction. It follows that $\mathrm{rk} \mathbf{M} = |S| - 1 = \mathrm{rk}(\mathbf{M} \setminus H) + |H| - 1$ and hence the second equality. \square

Proposition 2.5 (Handles in 3-connected matroids). *Let \mathbf{M} be a 3-connected matroid on E with $|E| > 3$. Then all its handles are non-disconnective 1-handles.*

Proof. Let $H \in \mathcal{H}_{\mathbf{M}}$ be any handle. By Lemma 2.4.(a), H must be proper. By Lemma 2.4.(e), H is not separating, that is, $|H| = 1$ or $|E \setminus H| = 1$. In the latter case, \mathbf{M} is a circuit by Lemma 2.4.(b) and hence not 3-connected (see Example 2.2). So H is a 1-handle.

Suppose that H is disconnective. Consider the deletion $\mathbf{M}' := \mathbf{M} \setminus H$ on the set $E' := E \setminus H$. Pick a connected component X of \mathbf{M}' of minimal size $|X| < |E'|$. Since $H \neq \emptyset$ and $|E| > 3$, both $X \cup H$ and its complement $E \setminus (X \cup H) = E' \setminus X$ have at least 2 elements. Since X is a connected component of \mathbf{M}' and by Lemma 2.4.(e),

$$\mathrm{rk}(X) + \mathrm{rk}(E' \setminus X) = \mathrm{rk} \mathbf{M}' = \mathrm{rk} \mathbf{M}.$$

Since $\mathrm{rk}(X \cup H) \leq \mathrm{rk}(X) + |H| = \mathrm{rk}(X) + 1$, it follows that

$$\lambda_{\mathbf{M}}(X \cup H) = \mathrm{rk}(X \cup H) + \mathrm{rk}(E \setminus (X \cup H)) - \mathrm{rk} \mathbf{M} < 2.$$

Whence $X \cup H$ is a 2-separation of \mathbf{M} , a contradiction. \square

The following notion is the basis for our inductive approach to connected matroids.

Definition 2.6 (Handle decompositions). Let M be a connected matroid. A *handle decomposition* of length k of M is a filtration

$$\mathcal{C}_M \ni F_1 \subsetneq \cdots \subsetneq F_k = E$$

such that $M|_{F_i}$ is connected and $H_i := F_i \setminus F_{i-1} \in \mathcal{H}_{M|_{F_i}}$ for $i = 2, \dots, k$.

By Lemma 2.4.(b) and (2.5), a handle decomposition yields circuits

$$(2.12) \quad C_1 := F_1 \in \mathcal{C}_M, \quad H_i \subsetneq C_i \in \mathcal{C}_{M|_{F_i}} \subseteq \mathcal{C}_M, \quad i = 2, \dots, k.$$

Conversely, it can be constructed from a suitable sequence of circuits.

Example 2.7 (Handle decomposition of the prism matroid). The prism matroid (see Example 2.1) has handle partition

$$E = \{e_1, e_2\} \sqcup \{e_3, e_4\} \sqcup \{e_5, e_6\}.$$

A handle decomposition of length 2 is given by

$$F_1 = \{e_1, e_2, e_3, e_4\} \subsetneq F_2 = E.$$

Note that all handles are proper, maximal, separating 2-handles. \diamond

Proposition 2.8 (Existence of handle decompositions). *Let M be a connected matroid and $C_1 \in \mathcal{C}_M$. Then there is a handle decomposition of M starting with $F_1 = C_1$.*

Proof. There is a sequence of circuits $C_1, \dots, C_k \in \mathcal{C}_M$ which defines a filtration $F_i := \bigcup_{j \leq i} C_j$ such that $C_i \cap F_{i-1} \neq \emptyset$ and $C_i \setminus F_{i-1} \in \mathcal{C}_{M/F_{i-1}}$ for $i = 2, \dots, k$ (see [CH96]). The hypothesis $C_i \cap F_{i-1} \neq \emptyset$ implies that $M|_{F_i}$ is connected for $i = 1, \dots, k$.

It remains to check that $H_i = C_i \setminus F_{i-1} \in \mathcal{H}_{M|_{F_i}}$ for $i = 2, \dots, k$. Since circuits are nonempty, $\emptyset \neq H_i \subsetneq F_i$. Let $C \in \mathcal{C}_{M|_{F_i}}$ be a circuit such that $e \in C \cap H_i \subseteq C \cap C_i$. Suppose by way of contradiction that $H_i \not\subseteq C$. Then there exists some $d \in C_i \setminus (C \cup F_{i-1})$. By the strong circuit elimination axiom (see [Oxl11, Prop. 1.4.12]), there is a circuit $C' \in \mathcal{C}_{M|_{F_i}} \subseteq \mathcal{C}_M$ (see (2.5)) for which $d \in C' \subseteq (C \cup C_i) \setminus \{e\}$. Then $C' \setminus F_{i-1} \subseteq C_i \setminus F_{i-1} \in \mathcal{C}_{M/F_{i-1}}$ by assumption on C_i . It follows that either $C' \subseteq F_{i-1}$ or $C' \setminus F_{i-1} = C_i \setminus F_{i-1}$ (see (2.7)). The former is impossible because $C' \ni d \notin F_{i-1}$, and the latter because $C' \cup F_{i-1} \not\subseteq e \in C_i$. \square

In the sequel, we develop a bound for the number of non-disconnective handles.

Lemma 2.9 (Non-disconnective handles). *Let M be a connected matroid. Suppose that $H \in \mathcal{H}_M$ and $H' \in \mathcal{H}_{M \setminus H}$ are non-disconnective with $H \cup H' \neq E$. Then there is a non-disconnective handle $H'' \in \mathcal{H}_M$ such that $H'' \subseteq H'$, with equality if $H' \in \mathcal{H}_M$.*

Proof. By hypothesis, M and $M \setminus H$ are connected and $H \cup H' \neq E$ implies that both H and H' are proper handles. Then Lemma 2.4.(b) yields circuits $C \in \mathcal{C}_M$ and $C' \in \mathcal{C}_{M \setminus H} \subseteq \mathcal{C}_M$ (see (2.5)) such that $H \subsetneq C$ and $H' \subsetneq C'$.

Suppose that $C \subseteq H \cup H'$. Then the strong circuit elimination axiom (see [Oxl11, Prop. 1.4.12]) yields a circuit $C'' \in \mathcal{C}_M$ for which $C'' \subseteq H \cup C'$, $H' \not\subseteq C''$ and $C'' \not\subseteq H \cup H'$. Since $C'' \subsetneq C'$ contradicts incomparability of circuits, $H \subsetneq C''$ since H is a handle and Lemma 2.4.(b) forbids equality.

Replacing C by C'' if necessary, we may assume that $H' \not\subseteq C$ and $C \not\subseteq H \cup H'$. In particular, $H'' := H' \setminus C \in \mathcal{H}_{M \setminus H}$ and $H'' = H'$ if $H' \in \mathcal{H}_M$. Since $M \setminus (H \cup H')$ is connected by hypothesis, C witnesses the fact that H , $C \cap H'$ and $E \setminus (H \cup H')$ are in the same connected component of $M \setminus H''$ (see (2.5)). In other words, $M \setminus H''$ is connected. If $H'' \in \mathcal{H}_M$ is a handle, then H'' is therefore non-disconnective.

Otherwise, there is a circuit $C'' \in \mathcal{C}_M$ such that $\emptyset \neq C'' \cap H'' \neq H''$. In particular, $H \subseteq C''$ since otherwise $C'' \cap H = \emptyset$ and $C'' \in \mathcal{C}_{M \setminus H}$ (see (2.5)) which would contradict $H'' \in \mathcal{H}_{M \setminus H}$. This means that C'' connects H with $C'' \cap H''$. We may therefore replace H'' by $\emptyset \neq H'' \setminus C'' \subsetneq H''$ and iterate. Then $H'' \in \mathcal{H}_M$ after finitely many steps. \square

Lemma 2.10 (Handle decomposition of length 2). *Let M be a connected matroid with a handle decomposition of length 2. Then M has at least 3 (disjoint) non-disconnective handles. In case of equality, they form the handle partition of M .*

Proof. Consider the circuits $C' := C_1 \in \mathcal{C}_M$, $C := C_2 \in \mathcal{C}_M$ (see (2.12)), the non-disconnective handle $H := H_2 \in \mathcal{H}_M$ and the subsets $\emptyset \neq H' := C' \setminus C \subseteq E$ and $\emptyset \neq H'' := C \cap C' \subseteq E$. Then $E = H \sqcup H' \sqcup H''$ and $C' = H' \cup H''$ and $C = H \cup H''$.

Let $C'' \in \mathcal{C}_M$ be any circuit with $C' \neq C'' \neq C$. By incomparability of circuits, $C'' \not\subseteq C'$ and hence $H \subseteq C''$ since H is a handle. By Lemma 2.4.(d), we may assume that $|H| = 1$. Then $H' \subseteq C''$ (see [Oxl11, §1.1, Exc. 5]). In particular, $H' \in \mathcal{H}_M$ is a third non-disconnective handle. If $H \cup H' \subseteq C''$ is an equality, then also $H'' \in \mathcal{H}_M$ is a non-disconnective handle and $H \sqcup H' \sqcup H''$ is the handle decomposition.

Otherwise, C'' witnesses the fact that H , H' and $\emptyset \neq C'' \cap H'' \neq H''$ are in the same connected component of $M|_{C''}$ (see (2.5)). If $H'' \setminus C'' \in \mathcal{H}_M$ is a handle, then it is therefore non-disconnective. Otherwise, iterating yields a third non-disconnective handle $H'' \setminus C'' \supseteq H''' \in \mathcal{H}_M$. \square

Example 2.11 (Unexpected handles). Consider the matroid M on $E = \{1, \dots, 6\}$ whose bases

$$\begin{aligned} \mathcal{B}_M = \{ & \{1, 2, 3, 4\}, \{1, 2, 3, 5\}, \{1, 2, 4, 5\}, \{1, 3, 4, 5\}, \{2, 3, 4, 5\}, \\ & \{1, 2, 3, 6\}, \{1, 2, 4, 6\}, \{1, 3, 4, 6\}, \{2, 3, 4, 6\}, \\ & \{1, 3, 5, 6\}, \{1, 4, 5, 6\}, \{2, 3, 5, 6\}, \{2, 4, 5, 6\} \} \end{aligned}$$

index those sets of columns of the matrix

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 2 \\ 0 & 0 & 0 & 1 & 1 & 2 \end{pmatrix}$$

which form a basis of \mathbb{F}_3^4 (see Remark 2.15). Its circuits and maximal handles are given by

$$\mathcal{C}_M = \{F_1 := \{1, 2, 3, 4, 5\}, \{1, 2, 3, 4, 6\}, \{1, 2, 5, 6\}, \{3, 4, 5, 6\}\},$$

$$\text{Max } \mathcal{H}_M = \{\{1, 2\}, \{3, 4\}, \{5\}, \{6\} =: H_2\}.$$

In particular, M is connected with a handle decomposition

$$F_1 \subsetneq F_1 \sqcup H_2 =: F_2 = E$$

of length 2. Here all 4 maximal handles are non-disconnective and the inequality in Lemma 2.10 is strict. This can happen because M is not a graphic matroid (see Lemma 2.25). \diamond

Proposition 2.12 (Lower bound for non-disconnective handles). *Let M be a connected matroid with a handle decomposition of length $k \geq 2$. Then M has at least $k + 1$ (disjoint) non-disconnective handles.*

Proof. We argue by induction on k . The base case $k = 2$ is covered by Lemma 2.10. Suppose now that $k \geq 3$. By hypothesis (see Definition 2.6), $H_k \in \mathcal{H}_M$ is a non-disconnective handle and the connected matroid $M \setminus H_k = M|_{F_{k-1}}$ has a handle decomposition of length $k - 1$. By induction, there are k (disjoint) non-disconnective handles $H'_0, \dots, H'_{k-1} \in \mathcal{H}_{M \setminus H_k}$. Since $k \geq 3$ and handles are non-empty, $H_k \cup H'_i \neq E$ for $i = 0, \dots, k - 1$. For each $i = 0, \dots, k - 1$, Lemma 2.9 now yields a non-disconnective handle $H'_i \supseteq H''_i \in \mathcal{H}_M$. Thus, M has $k + 1$ (disjoint) non-disconnective handles $H''_0, \dots, H''_{k-1}, H_k \in \mathcal{H}_M$. \square

We conclude this section with an observation.

Lemma 2.13 (Existence of circuits). *Let M be a connected matroid of rank $\text{rk } M \geq 2$. Then there is a circuit $C \in \mathcal{C}_M$ of size $|C| \geq 3$.*

Proof. Pick $e \in E$. Since M is connected, E is the union of all circuits $e \in C \in \mathcal{C}_M$. Suppose that there are only 2-circuits. Then $E = \text{cl}_M(e)$ (see [Oxl11, Prop. 1.4.11.(ii)]) and hence $\text{rk } M = 1$ (see (2.2)), a contradiction. \square

2.3. Configurations and realizations. Our objects of interest are not associated with a matroid itself but with a realization as defined in the following. All matroid operations we consider come with a counterpart for realizations. For graphic matroids, these agree with familiar operations on graphs (see §2.4).

Fix a field \mathbb{K} and denote the \mathbb{K} -dualizing functor by

$$-\vee := \text{Hom}_{\mathbb{K}}(-, \mathbb{K}).$$

We consider a finite set E as a basis of the *based* \mathbb{K} -vector space \mathbb{K}^E and denote by $E^\vee = (e^\vee)_{e \in E}$ the dual basis of

$$(2.13) \quad (\mathbb{K}^E)^\vee = \mathbb{K}^{E^\vee}.$$

By abuse of notation, we set $S^\vee := (e^\vee)_{e \in S}$ for any subset $S \subseteq E$.

We consider configurations as defined by Bloch, Esnault and Kreimer (see [BEK06, §1]).

Definition 2.14 (Configurations and realizations). Let E be a finite set. A \mathbb{K} -vector subspace $W \subseteq \mathbb{K}^E$ is called a *configuration* (over \mathbb{K}). It defines a matroid \mathbf{M}_W on E with independent sets

$$(2.14) \quad \mathcal{I}_{\mathbf{M}_W} = \{S \subseteq E \mid S^\vee|_W \text{ is } \mathbb{K}\text{-linearly independent in } W^\vee\}.$$

Let \mathbf{M} be a matroid and $W \subseteq \mathbb{K}^E$ a configuration (over \mathbb{K}). If $\mathbf{M} = \mathbf{M}_W$, then W is called a (*linear*) *realization* of \mathbf{M} and \mathbf{M} is called (*linearly*) *realizable* (over \mathbb{K}). A matroid is called *binary* if it is realizable over \mathbb{F}_2 . A configuration $W \subseteq \mathbb{K}^E$ is called *totally unimodular* if $\text{ch } \mathbb{K} = 0$ and W admits a basis whose coefficient matrix with respect to E has all (maximal) minors in $\{0, \pm 1\}$. A matroid is called *regular* if it admits a totally unimodular realization. Equivalently, a regular matroid is realizable over every field (see [Oxl11, Thm. 6.6.3]).

Since $E^\vee|_W$ generates W^\vee , we have (see (2.14))

$$(2.15) \quad \text{rk}(\mathbf{M}_W) = \dim W^\vee = \dim W.$$

Remark 2.15 (Matroids and linear algebra). The notions in matroid theory (see §2.1) are derived from linear (in)dependence over \mathbb{K} . Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid \mathbf{M} . Pick a basis $w = (w^1, \dots, w^r)$ of W where $r := \text{rk } \mathbf{M}$ (see (2.15)). For each $e \in E$, $e^\vee|_W$ is then represented by the vector $(w_e^i)_i \in \mathbb{K}^r$ where $w_e^i := e^\vee(w^i)$ for $i = 1, \dots, r$. Order $E = \{e_1, \dots, e_n\}$ and set $w_j^i := w_{e_j}^i$ for $j = 1, \dots, n$. Then these vectors form the columns of the coefficient matrix $A = (w_j^i)_{i,j} \in \mathbb{K}^{r \times n}$ of w . By construction, W is the row span of A . The matroid rank $\text{rk}_{\mathbf{M}}(S)$ of any subset $S \subseteq E$ now equals the \mathbb{K} -linear rank of the submatrix of A with columns S (see (2.1) and (2.14)). An element $e \in E$ is a loop in \mathbf{M} if and only if column e of A is zero; e is a coloop in \mathbf{M} if and only if column e is not in the span of the other columns. \diamond

Remark 2.16 (Classical configurations). Suppose that \mathbf{M}_W has no loops, that is, $e^\vee|_W \neq 0$ for each $e \in E$. Then the images of the $e^\vee|_W$ in $\mathbb{P}W^\vee$ form a *projective point configuration* in the classical sense (see [HC52]). Dually, the hyperplanes $\ker(e^\vee) \cap W$ form a *hyperplane arrangement* in W (see [OT92]), which is an equivalent notion in this case. \diamond

We fix some notation for realizations of basic matroid operations. Any subset $S \subseteq E$ gives rise to an inclusion and a projection

$$(2.16) \quad \iota_S: \mathbb{K}^S \hookrightarrow \mathbb{K}^E, \quad \pi_S: \mathbb{K}^E \twoheadrightarrow \mathbb{K}^E / \mathbb{K}^{E \setminus S} = \mathbb{K}^S$$

of based \mathbb{K} -vector spaces.

Definition 2.17 (Realizations of matroid operations). Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid \mathbf{M} , and let $F \subseteq E$ be any subset.

(a) The *restriction configuration* (see (2.16))

$$\begin{aligned} W|_F &:= \pi_F(W) \subseteq \mathbb{K}^F \\ &\cong (W + \mathbb{K}^{E \setminus F}) / \mathbb{K}^{E \setminus F} \cong W / (W \cap \mathbb{K}^{E \setminus F}) \end{aligned}$$

realizes the restriction matroid $\mathbf{M}|_F$.

(b) The *deletion configuration*

$$W \setminus F := W|_{E \setminus F}$$

realizes the deletion matroid $\mathbf{M} \setminus F$. We write $W \setminus e := W \setminus \{e\}$ for $e \in E$.

(c) The *contraction configuration*

$$W/F := W \cap \mathbb{K}^{E \setminus F} \subseteq \mathbb{K}^{E \setminus F}$$

realizes the contraction matroid \mathbf{M}/F .

(d) The *dual configuration* (see (2.13))

$$W^\perp := (\mathbb{K}^E / W)^\vee \subseteq \mathbb{K}^{E^\vee}$$

realizes the dual matroid \mathbf{M}^\perp .

(e) Any $0 \neq \varphi \in W^\vee$ defines an *elementary quotient configuration*

$$W_\varphi := \ker \varphi \subseteq \mathbb{K}^E.$$

Remark 2.18. Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid \mathbf{M} .

- (a) An element $e \in E$ is a loop or coloop in \mathbf{M} if and only if $W \subseteq \mathbb{K}^{E \setminus \{e\}}$ or $W = (W \setminus e) \oplus \mathbb{K}^{\{e\}}$, respectively. In both cases, $W \setminus e = W/e \subseteq \mathbb{K}^{E \setminus \{e\}}$.
- (b) For $0 \neq \varphi \in W^\vee$, pick $w \in W \setminus W_\varphi$ and $e \notin E$. Consider the configuration

$$W_{\varphi, w} := W_\varphi \oplus \mathbb{K} \cdot (w + e) \subseteq \mathbb{K}^{E \sqcup \{e\}}.$$

Then $W_{\varphi, w} \setminus e = W$ and $W_{\varphi, w}/e = W_\varphi$. By definition, \mathbf{M}_{W_φ} is therefore an *elementary quotient* of \mathbf{M}_W ; it can be characterized in terms of the notion of a *modular cut* (see [Kat16, §5.5] and [Oxl11, §7.3]). \diamond

Lemma 2.19 (Lift of direct sums to realizations). *Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid \mathbf{M} . Suppose that $\mathbf{M} = \mathbf{M}_1 \oplus \mathbf{M}_2$ decomposes with underlying partition $E = E_1 \sqcup E_2$. Then $W = W_1 \oplus W_2$ where $W_i := \mathbf{M}/E_j \subseteq \mathbb{K}^{E_i}$ realizes $\mathbf{M}_i = \mathbf{M}|_{E_i}$ for $\{i, j\} = \{1, 2\}$.*

Proof. By definition (see Definition 2.17.(a) and (c)),

$$W_1 \oplus W_2 \hookrightarrow W \hookrightarrow W|_{E_1} \oplus W|_{E_2}, \quad W_i \hookrightarrow W|_{E_i}, \quad i = 1, 2.$$

By the direct sum hypothesis, W_i and $W|_{E_i}$ realize the same matroid (see (2.3), (2.4) and (2.6))

$$\mathbf{M}/E_j = \mathbf{M}|_{E_i} = \mathbf{M}_i, \quad \{i, j\} = \{1, 2\}.$$

Thus, $\dim W_i = \dim(W|_{E_i})$ for $i = 1, 2$ (see (2.15)) and the claim follows. \square

Example 2.20 (Realizations of uniform matroids). Let $W \subseteq \mathbb{K}^E$ be the row span of a matrix $A \in \mathbb{K}^{r \times n}$ (see Remark 2.15). If A is generic in the sense that all maximal minors of A are nonzero, then W realizes the uniform matroid $\mathbf{U}_{r,n}$ (see Example 2.2). \diamond

2.4. Graphic matroids. Configurations arising from graphs are the most prominent examples for our results. In this subsection, we review this construction and discuss important examples such as prism, wheel and whirl matroids.

A *graph* $G = (V, E)$ is a pair of finite sets V of *vertices* and E of (unoriented) *edges* where each edge $e \in E$ is associated with a set of one or two vertices in V . This allows for multiple edges between pairs of vertices, and loops at vertices.


A graph G determines a *graphic matroid* \mathbf{M}_G on the edge set E . Its independent sets are the *forests* and its circuits the *simple cycles* in G . Any graphic matroid comes from a (non-unique) *connected* graph (see [Oxl11, Prop. 1.2.9]). Unless specified otherwise, we therefore assume that G is connected. Then the bases of \mathbf{M}_G are the *spanning trees* of G (see [Oxl11, p. 18]),

$$(2.17) \quad \mathcal{B}_{\mathbf{M}_G} = \mathcal{T}_G.$$

Remark 2.21 (Graph and matroid connectivity). A *vertex cut* of a graph $G = (V, E)$ is a subset of V whose removal (together with all incident edges) disconnects G . If G has at least one pair of distinct non-adjacent vertices, then G is called *k-connected* if k is the minimal size of a vertex cut. Otherwise, G is $(|V| - 1)$ -connected by definition. Suppose that $|V| \geq 3$. Then \mathbf{M}_G is (2-)connected if and only if G is 2-connected and loopless (see [Oxl11, Prop. 4.1.7]). Provided that $|E| \geq 4$, \mathbf{M}_G is 3-connected if and only if G is 3-connected and simple (see [Oxl11, Prop. 8.1.9]). \diamond

Example 2.22 (Prism matroid as graphic matroid). The prism matroid (see Definition 2.1) is associated with the (2, 2, 2)-theta graph in Figure 2. In particular it is 3-connected as witnessed by the minimal vertex cut $\{v_1, v_2, v_3\}$ (see Remark 2.21). \diamond

Graphic matroids have realizations derived from the edge-vertex incidence matrix of the graph (see [BEK06, §2]). A choice of orientation on the edge set E turns the graph G into a CW-complex. This gives

FIGURE 2. The $(2, 2, 2)$ -theta graph with a choice of orientation.


rise to an exact sequence

$$(2.18) \quad 0 \longrightarrow H_1 \longrightarrow \mathbb{K}^E \xrightarrow{\delta} \mathbb{K}^V \xrightarrow{\sigma} H_0 \longrightarrow 0$$

$$\begin{array}{ccccccc} & & \Downarrow & & \Downarrow & & \\ & & (s \rightarrow t) \mapsto t - s & & \mathbb{K} & & \\ & & & & \Downarrow & & \\ & & & & v \mapsto 1 & & \end{array}$$

where $H_\bullet := H_\bullet(G, \mathbb{K})$ denotes the graph homology of G over \mathbb{K} . The dual exact sequence

$$(2.19) \quad 0 \longleftarrow H^1 \longleftarrow \mathbb{K}^{E^\vee} \xleftarrow{\delta^\vee} \mathbb{K}^{V^\vee} \longleftarrow H^0 \longleftarrow 0$$

involves the graph cohomology $H^\bullet := H^\bullet(G, \mathbb{K})$ of G over \mathbb{K} .

Definition 2.23 (Graph configurations). We call the image

$$\mathbb{K}^{E^\vee} \supseteq W_G := \delta^\vee(\mathbb{K}^{V^\vee}) \xleftarrow[\cong]{\overline{\delta^\vee}} \ker(\sigma)^\vee$$

of δ^\vee the *graph configuration* of the graph G over \mathbb{K} . Note that it is independent of the orientation chosen to define δ in (2.18).

For any $S \subseteq E$, the sequence (2.18) induces a short exact sequence

$$0 \longrightarrow H_1 \cap \mathbb{K}^S \longrightarrow \mathbb{K}^S \longrightarrow W_G^\vee.$$

By definition of M_G and M_{W_G} (see Definition 2.14) and since H_1 is generated by indicator vectors of (simple) cycles, we have

$$S \in \mathcal{I}_{M_G} \iff H_1 \cap \mathbb{K}^S = 0 \iff S \in \mathcal{I}_{M_{W_G}},$$

which implies that

$$M_G = M_{W_G}.$$

The configuration W_G is totally unimodular if $\text{ch } \mathbb{K} = 0$ (see [Oxl11, Lem. 5.1.4]) which makes M_G a regular matroid. By construction, $W_G^\perp = H_1 \subseteq \mathbb{K}^E$ realizes the dual matroid M_G^\perp (see Definition 2.17.(d)).

Example 2.24 (Configuration of the $(2, 2, 2)$ -theta graph). With the orientation of the $(2, 2, 2)$ -theta graph G depicted in Figure 2, the map

δ^\vee in (2.19) is represented by the transpose of the matrix

$$\begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \\ -1 & 0 & -1 & 0 & -1 & 0 \end{pmatrix}.$$

Its rows generate the graph configuration W_G realizing the prism matroid (see Example 2.22). \diamond

Lemma 2.25 (Characterization of the prism matroid). *Let \mathbf{M} be a connected matroid on $E = \{e_1, \dots, e_6\}$ with $|E| = 6$ whose handle partition*

$$E = H_1 \sqcup H_2 \sqcup H_3, \quad H_1 = \{e_1, e_2\}, \quad H_2 = \{e_3, e_4\}, \quad H_3 = \{e_5, e_6\},$$

is made of 3 maximal 2-handles (see Example 2.7 and Lemma 2.10). Then \mathbf{M} is the prism matroid (see Definition 2.1). Up to scaling E , it has the unique realization $W \subseteq \mathbb{K}^E$ with basis

$$w^1 := e_1 + e_2, \quad w^2 := e_3 + e_4, \quad w^3 := e_5 + e_6, \quad w^4 := e_1 + e_3 + e_5,$$

the graph configuration of the (2, 2, 2)-theta graph (see Example 2.24).

Proof. Each circuit $C \in \mathcal{C}_{\mathbf{M}}$ is a (non-empty) disjoint union of H_1, H_2, H_3 (see Definition 2.3). By Lemma 2.4.(b), no H_i is a circuit, but each H_i is properly contained in one. By hypothesis, E is not a maximal handle and hence $E \notin \mathcal{C}_{\mathbf{M}}$. Up to renumbering H_1, H_2, H_3 , this yields circuits $H_2 \sqcup H_3$ and $H_1 \sqcup H_3$. By the strong circuit elimination axiom (see [Oxl11, Prop. 1.4.12]), there is a third circuit $H_1 \sqcup H_2$. Then

$$\mathcal{C}_{\mathbf{M}} = \{C_1, C_2, C_3\}, \quad C_1 = H_2 \sqcup H_3, \quad C_2 = H_1 \sqcup H_3, \quad C_3 = H_1 \sqcup H_2,$$

identifies with the circuits of the prism matroid. It follows that \mathbf{M} must be the prism matroid.

Let $W \subseteq \mathbb{K}^E$ be any realization of \mathbf{M} . Then $\dim W = \text{rk } \mathbf{M} = 4$ (see (2.15) and (2.17)). Pick a basis $w = (w^1, \dots, w^4)$ of W and denote by $A = (w_j^i)_{i,j}$ the coefficient matrix (see Remark 2.15). We may assume that columns 2, 4, 6, 5 of A form an identity matrix. Since C_1 and C_2 are circuits, $w_3^1 = 0 \neq w_3^2$ and $w_1^2 = 0 \neq w_1^1$. Thus,

$$A = \begin{pmatrix} * & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & * & 1 & 0 & 0 \\ * & 0 & * & 0 & 0 & 1 \\ * & 0 & * & 0 & 1 & 0 \end{pmatrix}.$$

Since C_3 is a circuit, suitably replacing $w^3, w^4 \in \langle w^3, w^4 \rangle$, reordering H_3 and scaling e_1, e_3 makes

$$A = \begin{pmatrix} * & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & * & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & * & 1 \\ 1 & 0 & 1 & 0 & 1 & 0 \end{pmatrix},$$

where $w_1^1, w_3^2, w_5^3 \neq 0$. Now suitably scaling first w^1, w^2, w^3 and then e_2, e_4, e_6 makes

$$A = \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 & 1 & 0 \end{pmatrix}.$$

Now $w = (w^1, \dots, w^4)$ is the desired basis. \square

The following classes of matroids play a distinguished role in connection with 3-connectedness.

Example 2.26 (Wheels and whirls). For $n \geq 2$, the *wheel graph* W_n in Figure 3 is obtained from an n -cycle, the “rim”, by adding an additional vertex and edges, the “spokes”, joining it to each vertex in the rim. There is a partition of the set of edges

$$E = S \sqcup R, \quad S = \{s_1, \dots, s_n\}, \quad R = \{r_1, \dots, r_n\},$$

into the set S of spokes and the set R of edges in the rim. The symmetry suggests to use a cyclic index set $\mathbb{Z}_n := \mathbb{Z}/n\mathbb{Z} = \{1, \dots, n\}$.




FIGURE 3. The wheel graph W_n .

For $n \geq 3$, the *wheel matroid* is the graphic matroid $W_n := M_{W_n}$ on E . For $n \geq 2$, the *whirl matroid* is the (non-graphic) matroid on E obtained from M_{W_n} by *relaxation* of the rim R , that is,

$$\mathcal{B}_{W^n} := \mathcal{B}_{M_{W_n}} \sqcup \{R\}.$$

In terms of circuits, this means that

$$\mathcal{C}_{W^n} = \mathcal{C}_{M_{W_n}} \setminus R \sqcup \{\{s\} \sqcup R \mid s \in S\}.$$

The matroids W_n and W^n are 3-connected (see [Oxl11, Exa. 8.4.3]) of rank

$$\text{rk } W_n = n = \text{rk } W^n.$$

For each $i \in \mathbb{Z}_n$, $\{s_i, r_i, s_{i+1}\}$ is a triangle and $\{r_i, r_{i+1}, s_{i+1}\}$ a triad. Conversely, this property enforces $M \in \{W_n, W^n\}$ for any connected matroid M on E (see [Sey80, (6.1)]). \diamond

We describe all realizations of wheels and whirls up to equivalence. In particular, we recover the well-known fact that whirls are not binary.

Lemma 2.27 (Realizations of wheels and whirls). *Let $W \subseteq \mathbb{K}^E$ be any realization of $M \in \{W_n, W^n\}$. Up to scaling $E = S \sqcup R$, W has a basis*

$$(2.20) \quad w^1 = s_1 + r_1 - t \cdot r_n, \quad w^i = s_i + r_i - r_{i-1}, \quad i = 2, \dots, n,$$

where $t = 1$ if $M = W_n$, and $t \in \mathbb{K} \setminus \{0, 1\}$ if $M = W^n$.

Proof. Since $S \in \mathcal{B}_M$, we may assume that the coefficients of s_j in w^i form an identity matrix, that is, $w_{s_j}^i = \delta_{i,j}$. The triangle $\{s_j, r_j, s_{j+1}\}$ then forces $w_{r_j}^j, w_{r_j}^{j+1} \neq 0$ and $w_{r_j}^i = 0$ for all $i \in \mathbb{Z}_n \setminus \{j, j+1\}$. Suitably scaling $r_1, w^2, r_2, w^3, \dots, r_{n-1}, w^n, s_1, \dots, s_n$ successively yields (2.20). The claim on t follows from $R \in \mathcal{C}_{W_n}$ and $R \in \mathcal{B}_{W^n}$, respectively. \square

3. CONFIGURATION POLYNOMIALS AND FORMS

In this section, we develop Bloch's strategy of putting graph polynomials into the context of configuration polynomials and configuration forms. We lay the foundation for an inductive proof of our main result using a handle decomposition. In the process, we generalize some known results on graph polynomials to configuration polynomials.

3.1. Configuration polynomials. To prepare the definition of configuration polynomials we introduce some notation.

Let $W \subseteq \mathbb{K}^E$ be a configuration, and let $S \subseteq E$ be any subset. Compose the associated inclusion map with π_S to a map (see (2.16))

$$(3.1) \quad \alpha_{W,S}: W \hookrightarrow \mathbb{K}^E \xrightarrow{\pi_S} \mathbb{K}^S.$$

Fix an isomorphism

$$(3.2) \quad c_W: \mathbb{K} \xrightarrow{\cong} \bigwedge^{\dim W} W$$

and set $c_0 := \text{id}_{\mathbb{K}}$ for the zero vector space. Any basis of W gives rise to such an isomorphism and any two such isomorphisms differ by a nonzero multiple $c \in \mathbb{K}^*$. Up to sign or ordering E , we identify

$$(3.3) \quad \bigwedge^{|S|} \mathbb{K}^S = \mathbb{K}, \quad \bigwedge_{s \in S} s \mapsto 1,$$

as based vector spaces. Suppose that $|S| = \dim W$. Then the determinant

$$(3.4) \quad \det \alpha_{W,S}: \mathbb{K} \xrightarrow{\cong} \bigwedge^{|S|} W \xrightarrow{\bigwedge^{|S|} \alpha_{W,S}} \bigwedge^{|S|} \mathbb{K}^S = \mathbb{K}$$

is defined up to sign. Its square

$$(3.5) \quad c_{W,S} := (\det \alpha_{W,S})^2 \in \mathbb{K}$$

is defined up to a factor c^2 for some $c \in \mathbb{K}^*$ independent of S . Note that $\det \alpha_{0,\emptyset} = \text{id}_{\mathbb{K}}$ and hence $c_{0,\emptyset} = 1$. By definition (see (2.14)),

$$(3.6) \quad c_{W,S} \neq 0 \iff S \in \mathcal{B}_{M_W}.$$

Remark 3.1 (Compatibility of coefficients with restriction). Let $W \subseteq \mathbb{K}^E$ be a configuration, and let $S \subseteq F \subseteq E$ with $|S| = \dim W$. Then the maps (3.1) for W and $W|_F$ form a commutative diagram

$$\begin{array}{ccccc} & & \alpha_{W,S} & & \\ & & \curvearrowright & & \\ W & \hookrightarrow & \mathbb{K}^E & \xrightarrow{\pi_S} & \mathbb{K}^S \\ & \downarrow \cong & \downarrow \pi_F & & \parallel \\ \pi_F|_W & & & & \\ W|_F & \hookrightarrow & \mathbb{K}^F & \xrightarrow{\pi_S} & \mathbb{K}^S \\ & & \alpha_{W|_F,S} & & \\ & & \curvearrowleft & & \end{array}$$

and hence $c_{W,S} = c^2 \cdot c_{W|_F,S}$ for some $c \in \mathbb{K}^*$ independent of S . \diamond

Consider the dual basis $E^\vee = (e^\vee)_{e \in E}$ of E as coordinates on \mathbb{K}^E ,

$$(3.7) \quad x_e := e^\vee, \quad \partial_e := \frac{\partial}{\partial x_e}, \quad e \in E.$$

Given an enumeration of $E = \{e_1, \dots, e_n\}$, we write

$$x_i := x_{e_i}, \quad \partial_i := \partial_{e_i}, \quad i = 1, \dots, n.$$

For any subset $S \subseteq E$, we set

$$(3.8) \quad x_S := (x_e)_{e \in S}, \quad x^S := \prod_{e \in S} x_e, \quad x := x_E.$$

Definition 3.2 (Configuration polynomials). Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid M . Then the *configuration polynomial* of W is (see (3.5))

$$\psi_W := \sum_{B \in \mathcal{B}_M} c_{W,B} \cdot x^B \in \mathbb{K}[x].$$

Remark 3.3 (Well-definedness of configuration polynomials). Any two isomorphisms c_W (see (3.2)) differ by a nonzero multiple $c \in \mathbb{K}^*$. Using the isomorphism $c \cdot c_W$ in place of c_W replaces ψ_W by $c^2 \cdot \psi_W$. In other words, ψ_W is well-defined up to a nonzero constant square factor. Whenever ψ_W occurs in a formula, we mean that the formula holds true for a suitable choice of such a factor. \diamond

Remark 3.4 (Equivalence of configuration polynomials). Dividing $e \in E$ by $c \in \mathbb{K}^*$ multiplies both $x_e = e^\vee$ (see Remark 2.16) and the identifications (3.3) with $e \in S$ by c . For each $e \in B \in \mathcal{B}_M$, this multiplies $c_{W,B}$ by c^2 and x^B by c . This is equivalent to substituting $c^3 \cdot x_e$ for x_e in ψ_W . Scaling E thus results in scaling x in ψ_W .

However, dropping the equality (3.7) and scaling $e \in E$ for fixed x_e replaces W in ψ_W by a projectively equivalent realization (see [Oxl11, §6.3]). If M is binary, then all realizations of M over \mathbb{K} are projectively equivalent (see [Oxl11, Prop. 6.6.5]). The corresponding configuration polynomials are geometrically equivalent in this case. In general, however, there are geometrically different configuration polynomials for fixed M and \mathbb{K} (see Example 5.3). \diamond

Remark 3.5 (Degree of configuration polynomials). Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid M . Then (see (2.15) and (3.6))

$$\deg \psi_W = \text{rk } M = \dim W.$$

In particular, $\psi_W \neq 0$, and $\psi_W = 1$ if and only if $\text{rk } M = 0$. By definition, ψ_W is independent of (divided by) x_e if and only if $e \in E$ is a (co)loop in M . \diamond

Remark 3.6 (Matroid polynomials and regularity). For any matroid M , not necessarily realizable, there is a *matroid (basis) polynomial*

$$\psi_M := \sum_{B \in \mathcal{B}_M} x^B.$$

If M is regular, then $\psi_W = \psi_M$ for any totally unimodular realization W of M over \mathbb{K} . Conversely, this equality for some realization W over \mathbb{K} with $\text{ch } \mathbb{K} = 0$ establishes regularity of M . For regular M , all configuration polynomials over \mathbb{K} are geometrically equivalent (see Remark 3.4). In general, however, ψ_W and ψ_M are geometrically different (see Example 5.2). \diamond

Example 3.7 (Configuration polynomials of uniform matroids). Let $W \subseteq \mathbb{K}^E$ be a realization of a uniform matroid $M = U_{r,n}$ (see Example 2.20).

(a) Suppose that $M = U_{n,n}$ is a free matroid. Then $E \in \mathcal{B}_M$ and

$$\psi_W = x^E$$

is the elementary symmetric polynomial of degree n in n variables.

(b) Suppose that $M = U_{n-1,n}$ is a circuit. Then $E \in \mathcal{C}_M$ and by Remark 3.1 and (a)

$$\psi_W = \sum_{e \in E} \psi_{W \setminus e}, \quad \psi_{W \setminus e} = x^{E \setminus \{e\}}.$$

A priori, substituting $x^{E \setminus \{e\}}$ for $\psi_{W \setminus e}$ in ψ_W is invalid (see Remark 3.3). However, this can be achieved as follows: Ordering $E = \{e_1, \dots, e_n\}$, W has a basis $w^i = e_i + c_i \cdot e_n$ with $c_i \in \mathbb{K}^*$ where $i = 1, \dots, n-1$. Scaling

first w^1, \dots, w^{n-1} and then e_1, \dots, e_{n-1} makes $c_1 = \dots = c_{n-1} = 1$. This turns ψ_W into

$$\psi_W = \sum_{e \in E} x^{E \setminus \{e\}},$$

the elementary symmetric polynomial of degree $n - 1$ in n variables.

(c) If $\mathbf{M} = \mathbf{U}_{n-2,n}$, then \mathbf{M} has $\binom{n}{n-2}$ bases, and ψ_W has $\binom{n}{n-2}$ monomials whose coefficients depend on the choice of W . For instance, the row span W of the matrix

$$\begin{pmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & -1 \end{pmatrix}$$

realizes $\mathbf{U}_{2,4}$ and

$$\psi_W = x_1x_2 + x_1x_3 + x_1x_4 + x_2x_3 + x_2x_4 + 4x_3x_4.$$

Realizations of $\mathbf{U}_{2,n}$ are treated in Example 5.4. \diamond

In the following, we put matroid connectivity in correspondence with irreducibility of configuration polynomials.

Proposition 3.8 (Connectedness and irreducibility). *Let \mathbf{M} be a matroid of rank $\text{rk } \mathbf{M} \geq 1$ with realization $W \subseteq \mathbb{K}^E$. Then \mathbf{M} is connected if and only if \mathbf{M} has no loops and ψ_W is irreducible. In particular, if $\mathbf{M} = \bigoplus_{i=1}^n \mathbf{M}_i$ with connected components \mathbf{M}_i and induced decomposition $W = \bigoplus_{i=1}^n W_i$ (see Lemma 2.19), then $\psi_W = \prod_{i=1}^n \psi_{W_i}$ where ψ_{W_i} is irreducible if $\text{rk } \mathbf{M}_i \geq 1$, and $\psi_{W_i} = 1$ otherwise.*

Proof. First suppose that $\mathbf{M} = \mathbf{M}_1 \oplus \mathbf{M}_2$ is disconnected with underlying proper partition $E = E_1 \sqcup E_2$. By Lemma 2.19, $W = W_1 \oplus W_2$ where $W_i \subseteq \mathbb{K}^{E_i}$ realizes \mathbf{M}_i . Then $\alpha_{W,B} = \alpha_{W_1,B_1} \oplus \alpha_{W_2,B_2}$ and hence $c_{W,B} = c_{W_1,B_1} \cdot c_{W_2,B_2}$ for all $B = B_1 \sqcup B_2 \in \mathcal{B}_{\mathbf{M}}$ where $B_i \in \mathcal{B}_{\mathbf{M}_i}$ for $i = 1, 2$ (see (2.3)). It follows that $\psi_W = \psi_{W_1} \cdot \psi_{W_2}$. This factorization is proper if \mathbf{M} and hence each \mathbf{M}_i has no loops (see Remark 3.5). Thus, ψ_W is reducible in this case.

Suppose now that ψ_W is reducible. Then

$$\psi_W = \psi_1 \cdot \psi_2$$

with ψ_i homogeneous non-constant for $i = 1, 2$. Since ψ_W is a linear combination of square-free monomials (see Definition 3.2), this yields a proper partition $E = E_1 \sqcup E_2$ such that $\psi_i \in \mathbb{K}[x_{E_i}]$ for $i = 1, 2$. Set

$$(3.9) \quad \mathbf{M}_i := \mathbf{M}|_{E_i}, \quad i = 1, 2.$$

Each basis $B \in \mathcal{B}_{\mathbf{M}}$ indexes a monomial x^B in ψ_W (see (3.6)). Set $B_i := B \cap E_i \in \mathcal{I}_{\mathbf{M}_i}$ for $i = 1, 2$ (see (2.4)). Then $x^B = x^{B_1} \cdot x^{B_2}$ where x^{B_i} is a monomial in ψ_i for $i = 1, 2$. By homogeneity of ψ_i , $B_i \in \mathcal{B}_{\mathbf{M}_i}$ for $i = 1, 2$ and hence $B = B_1 \sqcup B_2 \in \mathcal{B}_{\mathbf{M}_1 \oplus \mathbf{M}_2}$ (see (2.3)). It follows that $\mathcal{B}_{\mathbf{M}} \subseteq \mathcal{B}_{\mathbf{M}_1 \oplus \mathbf{M}_2}$.

Conversely, let $B = B_1 \sqcup B_2 \in \mathcal{B}_{\mathbf{M}_1 \oplus \mathbf{M}_2}$ where $B_i \in \mathcal{B}_{\mathbf{M}_i}$ for $i = 1, 2$. Then $B_i = B'_i \cap E_i$ for some $B'_i \in \mathcal{B}_{\mathbf{M}}$ for $i = 1, 2$ (see (2.4) and (3.9)).

As above, x^{B_i} is a monomial in ψ_i for $i = 1, 2$. Then $x^B = x^{B_1} \cdot x^{B_2}$ is a monomial in ψ_W and hence $B \in \mathcal{B}_M$ (see (3.6)). It follows that $\mathcal{B}_M \supseteq \mathcal{B}_{M_1 \oplus M_2}$ as well.

So $M = M_1 \oplus M_2$ is a proper decomposition and M is disconnected.

This proves the equivalence and the particular claims follow. \square

We use the following well-known fact from linear algebra.

Remark 3.9 (Determinant formula). Consider a short exact sequence of finite dimensional \mathbb{K} -vector spaces

$$0 \longrightarrow W \longrightarrow V \longrightarrow U \longrightarrow 0.$$

Abbreviate $\bigwedge V := \bigwedge^{\dim V} V$. There is a unique isomorphism

$$\bigwedge W \otimes \bigwedge U = \bigwedge V$$

that fits into a commutative diagram of canonical maps

$$\begin{array}{ccc} \bigwedge W \otimes \bigwedge^{\dim U} V & \longrightarrow & \bigwedge^{\dim W} V \otimes \bigwedge^{\dim U} V \\ \downarrow & & \downarrow \\ \bigwedge W \otimes \bigwedge U & \xlongequal{\quad} & \bigwedge V. \end{array}$$

Tensoring with

$$(\bigwedge U)^\vee = \bigwedge(U^\vee), \quad (\bigwedge W)^\vee = \bigwedge(W^\vee),$$

respectively, it induces identifications

$$\bigwedge W = \bigwedge V \otimes \bigwedge U^\vee, \quad \bigwedge U = \bigwedge W^\vee \otimes \bigwedge V.$$

Consider a commutative diagram of finite dimensional \mathbb{K} -vector spaces with short exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & W & \longrightarrow & V & \longrightarrow & U \longrightarrow 0 \\ & & \alpha \downarrow \cong & & \gamma \downarrow \cong & & \beta \uparrow \cong \\ 0 & \longleftarrow & U' & \longleftarrow & V' & \longleftarrow & W' \longleftarrow 0 \end{array}$$

Then the above identifications for both rows fit into a commutative diagram

$$\begin{array}{ccc} \bigwedge W & \xlongequal{\quad} & \bigwedge W \otimes \bigwedge U \otimes \bigwedge U^\vee & \xlongequal{\quad} & \bigwedge V \otimes \bigwedge U^\vee \\ \bigwedge \alpha \downarrow \cong & & \bigwedge \alpha \otimes \bigwedge \beta^{-1} \otimes \bigwedge \beta^\vee \downarrow \cong & & \bigwedge \gamma \otimes \bigwedge \beta^\vee \downarrow \cong \\ \bigwedge U' & \xlongequal{\quad} & \bigwedge U' \otimes \bigwedge W' \otimes \bigwedge W'^\vee & \xlongequal{\quad} & \bigwedge V' \otimes \bigwedge W'^\vee. \end{array}$$

\diamond

The following result of Bloch, Esnault and Kreimer describes the behavior of configuration polynomials under duality (see [BEK06, Prop. 1.6]).

Proposition 3.10 (Dual configuration polynomials). *Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid \mathbf{M} . For a suitable choice of c_W (see (3.2)),*

$$\det \alpha_{W^\perp, S^\perp} = \det \alpha_{W, S}$$

for all $S \subseteq E$ of size $|S| = \text{rk } \mathbf{M}$. In particular,

$$\psi_{W^\perp} = x^{E^\vee} \cdot \psi_W((x_{e^\vee}^{-1})_{e \in E}).$$

Proof. Let $S \subseteq E$ be of size $|S| = \text{rk } \mathbf{M}$. Then $S \in \mathcal{B}_\mathbf{M}$ if and only if $S^\perp \in \mathcal{B}_{\mathbf{M}^\perp}$ (see Remark 3.3). We may assume that this is the case as otherwise both determinants are zero. Then there is a commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & W & \longrightarrow & \mathbb{K}^E & \longrightarrow & \mathbb{K}^E/W \longrightarrow 0 \\ & & \alpha_{W,S} \downarrow \cong & & \nu \downarrow \cong & & \cong \uparrow \alpha_{W^\perp, S^\perp}^\vee \\ 0 & \longleftarrow & \mathbb{K}^S & \xleftarrow{\pi_{S^\perp} \circ \nu^{-1}} & \mathbb{K}^{E^\vee} & \xleftarrow{\pi_{S^\perp}^\vee} & \mathbb{K}^{S^\perp} \longleftarrow 0 \end{array}$$

where the middle isomorphism is induced by (2.8). This yields a commutative diagram (Remark 3.9 and (2.15))

$$\begin{array}{ccc} \mathbb{K} & \xrightarrow{\cong} & \bigwedge^{|E|} \mathbb{K}^E \otimes_{\mathbb{K}} \mathbb{K} \\ c_W \downarrow & & \downarrow \text{id} \otimes c_{W^\perp} \\ \bigwedge^{\text{rk } \mathbf{M}} W & \xlongequal{\quad} & \bigwedge^{|E|} \mathbb{K}^E \otimes \bigwedge^{\text{rk } \mathbf{M}^\perp} W^\perp \\ \bigwedge^{\text{rk } \mathbf{M}} \alpha_{W,S} \downarrow & & \downarrow \bigwedge^{|E|} \nu \otimes \bigwedge^{\text{rk } \mathbf{M}^\perp} \alpha_{W^\perp, S^\perp} \\ \bigwedge^{\text{rk } \mathbf{M}} \mathbb{K}^S & \xlongequal{\quad} & \bigwedge^{|E|} \mathbb{K}^{E^\vee} \otimes \bigwedge^{\text{rk } \mathbf{M}^\perp} \mathbb{K}^{S^\perp}. \end{array}$$

Using (3.3), we may drop $\bigwedge^{|E|} \mathbb{K}^E$ and $\bigwedge^{|E|} \mathbb{K}^{E^\vee}$. A suitable choice of c_W turns the upper isomorphism into an equality. The claim follows by definition (see (3.4) and Definition 3.2). \square

The coefficients of the configuration polynomial satisfy the following restriction–contraction formula.

Lemma 3.11 (Restriction–contraction for coefficients). *Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid \mathbf{M} , and let $F \subseteq E$ be any subset. For any basis $B \in \mathcal{B}_\mathbf{M}$, $B \cap F \in \mathcal{B}_{\mathbf{M}|_F}$ if and only if $B \setminus F \in \mathcal{B}_{\mathbf{M}/F}$. In this case,*

$$c_{W,B} = c^2 \cdot c_{W/F, B \setminus F} \cdot c_{W|_F, B \cap F}$$

where $c \in \mathbb{K}^*$ is independent of B .

Proof. The equivalence for $B \in \mathcal{B}_\mathbf{M}$ holds by definition of matroid contraction (see (2.6)). For any such B , there is a commutative diagram

with exact rows (see Definition 2.17.(a) and (c))

$$\begin{array}{ccccccc}
0 & \longrightarrow & W/F & \longrightarrow & W & \longrightarrow & W|_F \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \mathbb{K}^{E \setminus F} & \longrightarrow & \mathbb{K}^E & \longrightarrow & \mathbb{K}^F \longrightarrow 0 \\
& & \downarrow & & \downarrow & & \downarrow \\
0 & \longrightarrow & \mathbb{K}^{B \setminus F} & \longrightarrow & \mathbb{K}^B & \longrightarrow & \mathbb{K}^{B \cap F} \longrightarrow 0.
\end{array}$$

Taking exterior powers yields (see Remark 3.9 and (2.15))

$$\begin{array}{ccc}
\mathbb{K} & \xrightarrow[\cong]{c} & \mathbb{K} = \mathbb{K} \otimes \mathbb{K} \\
c_W \downarrow \cong & & \cong \downarrow c_{W/F} \otimes c_{W|_F} \\
\bigwedge^{\text{rk } M} W & \cong & \bigwedge^{\text{rk}(M/F)} W/F \otimes \bigwedge^{\text{rk}(M|_F)} W|_F \\
\bigwedge^{\text{rk } M} \alpha_{W,B} \downarrow \cong & & \cong \downarrow \bigwedge^{\text{rk}(M/F)} \alpha_{W/F, B \setminus F} \otimes \bigwedge^{\text{rk}(M|_F)} \alpha_{W|_F, B \cap F} \\
\bigwedge^{\text{rk } M} \mathbb{K}^B & \cong & \bigwedge^{\text{rk}(M/F)} \mathbb{K}^{B \setminus F} \otimes \bigwedge^{\text{rk}(M|_F)} \mathbb{K}^{B \cap F}.
\end{array}$$

□

The following result describes the behavior of configuration polynomials under deletion–contraction. It is the basis for our inductive approach to Jacobian schemes of configuration polynomials. The statement on $\partial_e \psi_W$ was proven by Patterson (see [Pat10, Lem. 4.4]).

Proposition 3.12 (Deletion–contraction for configuration polynomials). *Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid M , and let $e \in E$. Then*

$$\psi_W = \begin{cases} \psi_{W \setminus e} = \psi_{W/e} & \text{if } e \text{ is a loop in } M, \\ \psi_{W|_e} \cdot \psi_{W/e} = \psi_{W|_e} \cdot \psi_{W \setminus e} & \text{if } e \text{ is a coloop in } M, \\ \psi_{W \setminus e} + \psi_{W|_e} \cdot \psi_{W/e} & \text{otherwise,} \end{cases}$$

where $\psi_{W|_e} = x_e$ if e is not a loop in M . In particular,

$$\partial_e \psi_W = \begin{cases} 0 & \text{if } e \text{ is a loop in } M, \\ \psi_{W/e} = \psi_{W \setminus e} & \text{if } e \text{ is a coloop in } M, \\ \psi_{W/e} & \text{otherwise,} \end{cases}$$

$$\psi_W|_{x_e=0} = \begin{cases} \psi_{W \setminus e} = \psi_{W/e} & \text{if } e \text{ is a loop in } M, \\ 0 & \text{if } e \text{ is a coloop in } M, \\ \psi_{W \setminus e} & \text{otherwise.} \end{cases}$$

Proof. Decompose

$$(3.10) \quad \psi_W = \sum_{e \notin B \in \mathcal{B}_M} c_{W,B} \cdot x^B + x_e \cdot \sum_{e \in B \in \mathcal{B}_M} c_{W,B} \cdot x^{B \setminus \{e\}}.$$

The second sum in (3.10) is nonzero if and only if e is not a loop. Suppose that this is the case. Then $\mathbf{M}|_e$ is free with basis $\{e\}$ and $\psi_{W|_e} = x_e$ by Remark 3.7.(a). By Lemma 3.11 applied to $F = \{e\}$, the second sum in (3.10) then equals (see (2.6) and Remark 3.3)

$$c^2 \cdot c_{W|_e, \{e\}} \cdot \sum_{B \in \mathcal{B}_{\mathbf{M}|_e}} c_{W/e, B} \cdot x^B = \psi_{W/e}$$

for some $c \in \mathbb{K}^*$. The first sum in (3.10) is nonzero if and only if e is not a coloop. By Lemma 3.11 applied to $F = E \setminus \{e\}$, it equals in this case (see (2.4) and Remark 3.3)

$$c^2 \cdot c_{0, \emptyset} \cdot \sum_{B \in \mathcal{B}_{\mathbf{M} \setminus e}} c_{W \setminus e, B} \cdot x^B = \psi_{W \setminus e}$$

for some $c \in \mathbb{K}^*$. If e is a (co)loop, then $W/e = W \setminus e$ (see Remark 2.18.(a)). The claimed formulas follow. \square

The following formula relates configuration polynomials with deletion and contraction of handles. It is the starting point for our description of generic points of Jacobian schemes of configuration hypersurfaces in terms of handles.

Corollary 3.13 (Configuration polynomials and handles). *Let $W \subseteq \mathbb{K}^E$ be a realization of a connected matroid \mathbf{M} on E , and let $E \neq H \in \mathcal{H}_{\mathbf{M}}$ be a proper handle. Then*

$$(3.11) \quad \psi_W = \psi_{W/(E \setminus H)} \cdot \psi_{W \setminus H} + \psi_{W|_H} \cdot \psi_{W/H},$$

$$(3.12) \quad \psi_{W/(E \setminus H)} = \sum_{h \in H} \psi_{W|_{H \setminus \{h\}}},$$

$$(3.13) \quad \psi_{W|_H} = x^H, \quad \psi_{W|_{H \setminus \{h\}}} = x^{H \setminus \{h\}}.$$

In particular, after suitably scaling H ,

$$(3.14) \quad \psi_W = \sum_{h \in H} x^{H \setminus \{h\}} \cdot \psi_{W \setminus H} + x^H \cdot \psi_{W/H}.$$

Proof. By Lemma 2.4.(b), $H \in \mathcal{C}_{\mathbf{M}/(E \setminus H)}$ and hence (3.12) by Example 3.7.(b). By Lemma 2.4.(b) (see (2.4)), $\mathbf{M}|_H$ is free, and equalities (3.13) follows from Example 3.7.(a). Equality (3.14) follows from (3.11), (3.12) and Example 3.7.(b). It remains to prove equality (3.11).

We proceed by induction on $|H|$. Let $h \in H$ and set $H' := H \setminus \{h\}$. Since \mathbf{M} is connected, it has no (co)loops and hence

$$(3.15) \quad \psi_W = \psi_{W \setminus h} + \psi_{W|_h} \cdot \psi_{W/h}$$

by Proposition 3.12. If $|H| = 1$, then $H \in \mathcal{C}_{\mathbf{M}/(E \setminus H)}$ implies that $\text{rk}(\mathbf{M}/(E \setminus h)) = 0$ and hence $\psi_{W/(E \setminus h)} = 1$ (see Remark 3.5). Suppose now that $|H| \geq 2$. By Lemma 2.4.(b) and (c), $\mathbf{M}|_{H'}$ is free and H' consists of coloops in $\mathbf{M} \setminus h$. Iterating Proposition 3.12 thus yields

$$(3.16) \quad \psi_{W \setminus h} = \prod_{h' \in H'} \psi_{W|_{h'}} \cdot \psi_{W \setminus H} = \psi_{W|_{H'}} \cdot \psi_{W \setminus H}.$$

By Lemma 2.4.(d), the set H' is a proper handle in the connected matroid M/h . By Lemma 2.4.(c), h is a coloop in $M \setminus H'$ and hence

$$W/h \setminus H' = W \setminus H'/h = W \setminus H' \setminus h = W \setminus H.$$

by Remark 2.18.(a). By the induction hypothesis,

$$(3.17) \quad \psi_{W/h} = \sum_{h' \in H'} \psi_{W|_{H' \setminus \{h'\}}} \cdot \psi_{W \setminus H} + \psi_{W|_{H'}} \cdot \psi_{W/H}.$$

By Lemma 2.4.(b), $M|_H$ and $M|_{H \setminus \{h'\}}$ are free. Iterating Proposition 3.12 thus yields

$$(3.18) \quad \psi_{W|_h} \cdot \psi_{W|_{H'}} = \psi_{W|_H}, \quad \psi_{W|_h} \cdot \psi_{W|_{H' \setminus \{h'\}}} = \psi_{W|_{H \setminus \{h'\}}}.$$

Using equalities (3.12) and (3.18), equality (3.11) is obtained by substituting (3.16) and (3.17) into (3.15) (see Remark 3.3). \square

The following result describes the behavior of configuration polynomials when passing to an elementary quotient.

Proposition 3.14 (Configuration polynomials of quotients). *Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid M , and let $0 \neq \varphi \in W^\vee$. Then*

$$\psi_{W_\varphi} = \sum_{\substack{S \subseteq E \\ |S| = \text{rk } M - 1}} \left(\sum_{e \notin S} \pm \tilde{\varphi}_e \cdot \det \alpha_{W, S \cup \{e\}} \right)^2 x^S,$$

where $\tilde{\varphi} = (\tilde{\varphi}_e)_{e \in E} \in (\mathbb{K}^E)^\vee$ is any lift of φ with a sign \pm determined by a Laplace expansion.

Proof. Set $V := W^\perp$ and $V_\varphi := W_\varphi^\perp$ and consider the commutative diagram with short exact rows and columns

$$\begin{array}{ccccccc} & & & & 0 & & \\ & & & & \downarrow & & \\ & & & & \mathbb{K} & & \\ & & & & \downarrow & & \\ 0 & \longrightarrow & W_\varphi & \longrightarrow & \mathbb{K}^E & \longrightarrow & V_\varphi^\vee \longrightarrow 0 \\ & & \downarrow & & \parallel & & \downarrow \\ 0 & \longrightarrow & W & \longrightarrow & \mathbb{K}^E & \longrightarrow & V^\vee \longrightarrow 0 \\ & & \downarrow \varphi & \swarrow \tilde{\varphi} & & & \downarrow \\ & & \mathbb{K} & & & & 0 \\ & & \downarrow & & & & \\ & & 0 & & & & \end{array}$$

Dualizing and identifying the two copies of \mathbb{K} by the Snake Lemma yields a commutative diagram with short exact rows and columns

$$(3.19) \quad \begin{array}{ccccccc} & & & & & & 0 \\ & & & & & & \uparrow \\ & & & & & & \mathbb{K} \\ & & & & \cdot\tilde{\varphi} & & \uparrow \\ & & & & \mathbb{K}^{E^\vee} & \longleftarrow & V_\varphi \longleftarrow 0 \\ & & & & \parallel & & \uparrow \\ 0 & \longleftarrow & W_\varphi^\vee & \longleftarrow & \mathbb{K}^{E^\vee} & \longleftarrow & V \longleftarrow 0 \\ & & \uparrow & & \parallel & & \uparrow \\ & & W^\vee & \longleftarrow & \mathbb{K}^{E^\vee} & \longleftarrow & V \longleftarrow 0 \\ & & \uparrow & & \parallel & & \uparrow \\ & & \mathbb{K} & \xrightarrow{\cdot\tilde{\varphi}} & \mathbb{K}^{E^\vee} & & 0 \\ & & \uparrow & & \uparrow & & \\ & & 0 & & & & \end{array}$$

By Remark 3.9 and with a suitable choice of c_V (see Remark 3.3), the right vertical short exact sequence in (3.19) gives rise to a commutative square

$$\begin{array}{ccc} \mathbb{K} & \xrightarrow{c_{V_\varphi}} & \bigwedge^{\text{rk } \mathbf{M}^\perp + 1} V_\varphi \\ \parallel & & \parallel \\ \mathbb{K} & \xrightarrow{c_V} & \bigwedge^{\text{rk } \mathbf{M}^\perp} V \end{array}$$

Let $S' \subseteq E^\vee$ with $|S'| = \dim V_\varphi = \text{rk } \mathbf{M}^\perp + 1$ and denote (see (2.8))

$$\tilde{\varphi}_{S'} = (\tilde{\varphi}_{\nu^{-1}(e)})_{e \in S'} \in \mathbb{K}^{S'}.$$

Due to (3.19) the maps $\alpha_{V_\varphi, S'}$ (see (3.1)) and

$$(\tilde{\varphi}_{S'} \quad \alpha_{V_\varphi, S'}) : \mathbb{K} \oplus V \longrightarrow \mathbb{K}^{E^\vee} \xrightarrow{\pi_{S'}} \mathbb{K}^{S'}$$

agree after applying $\bigwedge^{\text{rk } \mathbf{M}^\perp + 1}$. Laplace expansion thus yields

$$\det \alpha_{V_\varphi, S'} = \sum_{e \in S'} \pm \tilde{\varphi}_{\nu^{-1}(e)} \cdot \det \alpha_{V, S' \setminus \{e\}}.$$

Let $S \subseteq E$ with $|S| = \dim W_\varphi = \text{rk } \mathbf{M} - 1$ and $S' = S^\perp$. Then Proposition 3.10 yields

$$c_{W_\varphi, S} = \left(\sum_{e \notin S} \pm \tilde{\varphi}_e \cdot \det \alpha_{W, S \cup \{e\}} \right)^2. \quad \square$$

3.2. Graph polynomials. We continue the discussion of graphic matroids from §2.4 and consider their configuration polynomials.

Definition 3.15 (Graph polynomials). The *(first) Kirchhoff polynomial* of a graph G over \mathbb{K} is the polynomial

$$\psi_G := \sum_{T \in \mathcal{T}_G} x^T \in \mathbb{K}[x].$$

Replacing x^T by $x^{E \setminus T}$ defines the *(first) Symanzik polynomial* ψ_G^\perp of a graph G over \mathbb{K} . We refer to ψ_G and ψ_G^\perp as *(first) graph polynomials*.

By (2.17), we have $\psi_G = \psi_W$ for any totally unimodular realization W of M_G . In particular, this yields the following result of Bloch, Esnault and Kreimer (see [BEK06, Prop. 2.2] and Proposition 3.10).

Proposition 3.16 (Graph polynomials as configuration polynomials). *The graph polynomials*

$$\psi_G = \psi_{W_G}, \quad \psi_G^\perp = \psi_{W_G^\perp},$$

are the configuration polynomials of the graph configuration and of its dual (see Definition 2.23). \square

Example 3.17 (Graph polynomial of the prism). For the unique realization $W = W_G$ of the prism matroid (see Lemma 2.25),

$$\begin{aligned} \psi_W = \psi_G &= x_1 x_2 (x_3 + x_4) (x_5 + x_6) \\ &\quad + x_3 x_4 (x_1 + x_2) (x_5 + x_6) \\ &\quad + x_5 x_6 (x_1 + x_2) (x_3 + x_4) \end{aligned}$$

is the Kirchhoff polynomial of the $(2, 2, 2)$ -theta graph G (see Figure 2). \diamond

Let $G = (E, V)$ be a graph. A *2-forest* in G is an acyclic subgraph T of G with $|V| - 2$ edges. Any such $T = \{T_1, T_2\}$ has 2 connected components T_1 and T_2 . We denote by \mathcal{T}_G^2 the set of all 2-forests in G .

Definition 3.18 (Second graph polynomials). The *second Kirchhoff polynomial* of a graph G over \mathbb{K} is the polynomial

$$\psi_G(p) := \sum_{\{T_1, T_2\} \in \mathcal{T}_G^2} m_{T_1}(p)^2 \cdot x^{T_1 \sqcup T_2} \in \mathbb{K}[x], \quad m_{T_i}(p) := \sum_{v \in T_i} p_v,$$

depending on a *momentum* $0 \neq p \in \ker \sigma$ for G over \mathbb{K} (see (2.18)). Note that

$$m_{T_1}(p) = \sum_{v \in T_1} p_v = - \sum_{v \in T_2} p_v = -m_{T_2}(p),$$

and hence, the coefficient $m_{T_1}(p)^2 \in \mathbb{K}$ of $\psi_G(p)$ is well-defined.

Replacing the 2-forests $T_1 \sqcup T_2$ by *cut sets* $E \setminus (T_1 \sqcup T_2)$ defines the *second Symanzik polynomial* $\psi_G^\perp(p)$ of a graph G over \mathbb{K} (see [Pat10, Def. 3.6]). We refer to $\psi_G(p)$ and $\psi_G^\perp(p)$ as *second graph polynomials*.

The following reformulation of a result of Patterson realizes second graph polynomials as configuration polynomials of a (dual) elementary quotient (see [Pat10, Prop. 3.3] and Proposition 3.10). Patterson's proof makes the general formula in Proposition 3.14 explicit in case of graph configurations (see [Pat10, Lem. 3.4]).

Proposition 3.19 (Second graph polynomials as configuration polynomials). *The second graph polynomials*

$$\psi_G(p) = \psi_{(W_G)_p}, \quad \psi_G^\perp(p) = \psi_{((W_G)_p)^\perp},$$

are the configuration polynomials of the quotient of the graph configuration by a momentum and of its dual (see Definitions 2.17.(d) and (e) and 2.23). \square

3.3. Configuration forms. The configuration form yields an equivalent definition of the configuration polynomial as a determinant of a symmetric matrix with linear entries. Its second degeneracy locus turns out to be the non-smooth locus of the hypersurface defined by the corresponding configuration polynomial.

Definition 3.20 (Configuration forms). Let $\mu_{\mathbb{K}}$ denote the multiplication map of \mathbb{K} . Consider the generic diagonal bilinear form on \mathbb{K}^E ,

$$Q_{\mathbb{K}^E} := \sum_{e \in E} x_e \cdot \mu_{\mathbb{K}} \circ (e^\vee \times e^\vee): \mathbb{K}^E \times \mathbb{K}^E \rightarrow \mathbb{K}[x].$$

Let $W \subseteq \mathbb{K}^E$ be a configuration. Then the *configuration (bilinear) form* of W is the restriction of $Q_{\mathbb{K}^E}$ to W ,

$$Q_W := Q_{\mathbb{K}^E}|_{W \times W}: W \times W \rightarrow \mathbb{K}[x].$$

Alternatively, it can be seen as the composition of canonical maps

$$(3.20) \quad Q_W: W[x] \longrightarrow \mathbb{K}^E[x] \xrightarrow{Q_{\mathbb{K}^E}} \mathbb{K}^{E^\vee}[x] \longrightarrow W^\vee[x],$$

where $-[x]$ means $- \otimes \mathbb{K}[x]$. For $k = 0, \dots, r := \dim W$, it defines a map

$$\bigwedge^{r-k} W \otimes \bigwedge^{r-k} W \otimes \mathbb{K}[x] \rightarrow \mathbb{K}[x].$$

Its image is the k th Fitting ideal $\text{Fitt}_k \text{coker } Q_W$ (see [Eis95, §20.2]) and defines the $k - 1$ st degeneracy scheme of Q_W . We set

$$M_W := \text{Fitt}_1 \text{coker } Q_W \trianglelefteq \mathbb{K}[x].$$

Note the different fonts used for M_W and \mathbb{M}_W (see Definition 2.14).

Remark 3.21 (Configuration forms as matrices). With respect to a basis $w = (w^1, \dots, w^r)$ of W , Q_W becomes a matrix of Hadamard products (see Remark 2.15)

$$Q_w = (\langle x, w^i \star w^j \rangle)_{i,j} = \left(\sum_{e \in E} x_e \cdot w_e^i \cdot w_e^j \right)_{i,j} \in \mathbb{K}^{r \times r}, \quad w_e^i = e^\vee(w^i).$$

Let $Q^{i,j}$ denote the submaximal minor of a square matrix Q obtained by deleting row i and column j . Then

$$M_W = \langle Q_W^{i,j} \mid i, j \in \{1, \dots, r\} \rangle.$$

Any basis of W can be written as $w' = Uw$ for some $U \in \text{Aut}_{\mathbb{K}} W$. Then

$$Q_{w'} = UQ_wU^t.$$

and the $Q_{w'}^{i,j}$ become \mathbb{K} -linear combinations of the $Q_w^{i,j}$. We often consider Q_W as a matrix Q_w determined up to conjugation. \diamond

Remark 3.22 (Configuration forms and basis scaling). Scaling E results in scaling x in Q_W and in M_W (see Remark 3.4). \diamond

Bloch, Esnault and Kreimer defined ψ_W in terms of Q_W (see [BEK06, Lem. 1.3]).

Lemma 3.23 (Configuration polynomial from configuration form). *For any configuration $W \subseteq \mathbb{K}^E$, the configuration polynomial*

$$\psi_W = \det Q_W \in M_W$$

is the determinant of the configuration form (see Remarks 3.3 and 3.21). \square

Example 3.24 (Configuration form of the prism realization). Consider the realization W of the prism matroid with basis given in Lemma 2.25. Then the corresponding matrix of Q_W reads (see Remark 3.21)

$$Q_W = \begin{pmatrix} x_1 + x_2 & 0 & 0 & x_1 \\ 0 & x_3 + x_4 & 0 & x_3 \\ 0 & 0 & x_5 + x_6 & x_5 \\ x_1 & x_3 & x_5 & x_1 + x_3 + x_5 \end{pmatrix}.$$

Lemma 3.23 recovers the polynomial $\det Q_W = \psi_W$ in Example 3.17. \diamond

The following result describes the behavior of Fitting ideals of configuration forms under duality. We consider the torus

$$\mathbb{T}^E := (\mathbb{K}^*)^E \subset \mathbb{K}^E, \quad \mathbb{K}[\mathbb{T}^E] = \mathbb{K}[x^{\pm 1}] = \mathbb{K}[x]_{x^E}.$$

The *Cremona isomorphism* $\mathbb{T}^E \cong \mathbb{T}^{E^\vee}$ is defined by

$$(3.21) \quad \zeta_E: \mathbb{K}[\mathbb{T}^E] \cong \mathbb{K}[\mathbb{T}^{E^\vee}], \quad x_e^{-1} \leftrightarrow x_{e^\vee}, \quad e \in E.$$

Proposition 3.25 (Duality and cokernels of configuration forms). *Let $W \subseteq \mathbb{K}^E$ be a configuration. Then there is an isomorphism over ζ_E ,*

$$\text{coker}(Q_W)_{x^E} \cong \text{coker}(Q_{W^\perp})_{x^{E^\vee}},$$

where the indices denote localization (see (3.8)). In particular, this induces an isomorphism

$$(M_W)_{x^E} \cong (M_{W^\perp})_{x^{E^\vee}}.$$

Proof. Consider the short exact sequence

$$(3.22) \quad 0 \longrightarrow W \longrightarrow \mathbb{K}^E \longrightarrow \mathbb{K}^E/W \longrightarrow 0$$

and its \mathbb{K} -dual

$$(3.23) \quad 0 \longleftarrow W^\vee \longleftarrow \mathbb{K}^{E^\vee} \longleftarrow W^\perp \longleftarrow 0.$$

We identify $\mathbb{K}^E = \mathbb{K}^{E^{\vee\vee}}$ and $\mathbb{K}^E/W = W^{\perp\vee}$, and we abbreviate

$$Q := Q_{\mathbb{K}^E}, \quad Q^\vee := Q_{\mathbb{K}^{E^\vee}}.$$

Then Q_{x^E} and $Q_{x^{E^\vee}}^\vee$ are mutual inverses under ζ_E . Together with (3.22) and (3.23) tensored by $\mathbb{K}[x^{\pm 1}]$ and (3.20) for W and W^\perp , they fit into a commutative diagram with exact rows connected vertically by morphisms over ζ_E

$$\begin{array}{ccccccc}
 & & & & & & 0 \\
 & & & & & & \uparrow \\
 & & & & & & \text{coker}(Q_{W^\perp})_{x^{E^\vee}} \\
 & & & & & \nearrow & \uparrow \\
 0 & \longrightarrow & W[x^{\pm 1}] & \longrightarrow & \mathbb{K}^E[x^{\pm 1}] & \longrightarrow & W^{\perp\vee}[x^{\pm 1}] \longrightarrow 0 \\
 & & \downarrow (Q_W)_{x^E} & & \downarrow Q_{x^E} \uparrow Q_{x^{E^\vee}}^\vee & & \uparrow (Q_{W^\perp})_{x^{E^\vee}} \\
 0 & \longleftarrow & W^\vee[x^{\pm 1}] & \longleftarrow & \mathbb{K}^{E^\vee}[x^{\pm 1}] & \longleftarrow & W^\perp[x^{\pm 1}] \longleftarrow 0 \\
 & & \downarrow & & \nwarrow & & \uparrow \\
 & & \text{coker}(Q_W)_{x^E} & & & & 0 \\
 & & \downarrow & & & & \\
 & & 0, & & & &
 \end{array}$$

where $-[x^{\pm 1}]$ means $- \otimes \mathbb{K}[x^{\pm 1}]$. Exactness of the columns is due to $\det Q_W = \psi_W \neq 0$ (see Lemma 3.23 and Remark 3.5). Composing the middle vertical isomorphism over ζ_E with (taking preimages along) the dashed compositions yields the claimed isomorphism by a diagram chase. \square

The following result describes the behavior of submaximal minors of configuration forms under deletion–contraction. It is the basis for our inductive approach to second degeneracy schemes.

Lemma 3.26 (Deletion–contraction for submaximal minors). *Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid \mathbf{M} of rank $r = \text{rk } \mathbf{M}$, and let $e \in E$. Then any basis of W/e can be extended to bases of W and $W \setminus e$ such*

that $Q_W^{i,j} =$

$$\begin{cases} Q_{W \setminus e}^{i,j} = Q_{W/e}^{i,j} & \text{if } e \text{ is a loop in } \mathbf{M}, \\ \psi_{W \setminus e} = \psi_{W/e} & \text{if } e \text{ is a coloop in } \mathbf{M}, i = r = j, \\ x_e \cdot Q_{W \setminus e}^{i,j} = x_e \cdot Q_{W/e}^{i,j} & \text{if } e \text{ is a coloop in } \mathbf{M}, i \neq r \neq j, \\ 0 & \text{if } e \text{ is a coloop in } \mathbf{M}, \text{ otherwise,} \\ \psi_{W/e} & \text{if } e \text{ is not a (co)loop in } \mathbf{M}, i = r = j, \\ Q_{W \setminus e}^{i,j} & \text{if } e \text{ is not a (co)loop in } \mathbf{M}, i = r \text{ or } j = r, \\ Q_{W \setminus e}^{i,j} + x_e \cdot Q_{W/e}^{i,j} & \text{if } e \text{ is not a (co)loop in } \mathbf{M}, i \neq r \neq j, \end{cases}$$

for all $i, j \in \{1, \dots, r\}$. In particular, the $Q_W^{i,j}$ are linear combinations of square-free monomials for any basis of W .

Proof. Pick a basis w^1, \dots, w^r of $W \subseteq \mathbb{K}^E$ and consider

$$Q_W = \left(\sum_{e \in E} x_e \cdot w_e^i \cdot w_e^j \right)_{i,j} \in \mathbb{K}^{r \times r}$$

as a matrix (see Remark 3.21). Recall that (see Definition 2.17.(b) and (c)),

$$W \setminus e = \pi_{E \setminus \{e\}}(W), \quad W/e = W \cap \mathbb{K}^{E \setminus \{e\}},$$

and the description of (co)loops in Remark 2.18.(a):

- If e is a loop, then $w_e^i = 0$ for all $i = 1, \dots, r$ and hence $W \setminus e = W = W/e$.
- If e is not a loop, then we may adjust w^1, \dots, w^r such that $w_e^i = \delta_{i,r}$ for all $i = 1, \dots, r$ and then w^1, \dots, w^{r-1} is a general basis of W/e .
- If e is a coloop, then we may adjust $w^r = e$ and $\pi_{E \setminus \{e\}}$ identifies w^1, \dots, w^{r-1} with a basis of $W \setminus e = W/e$.

In the latter case,

$$(3.24) \quad Q_W = \begin{pmatrix} Q_{W \setminus e} & 0 \\ 0 & x_e \end{pmatrix},$$

and the claimed equalities follow (see Lemma 3.23).

It remains to consider the case in which e is not a (co)loop. Then $\iota_{E \setminus \{e\}}$ and $\pi_{E \setminus \{e\}}$ (see (2.16)) identify w^1, \dots, w^{r-1} and w^1, \dots, w^r with bases of W/e and $W \setminus e$, respectively. Hence,

$$(3.25) \quad Q_{W \setminus e} = \begin{pmatrix} Q_{W/e} & b \\ b^t & a \end{pmatrix}, \quad Q_W = \begin{pmatrix} Q_{W/e} & b \\ b^t & x_e + a \end{pmatrix},$$

where both the entry a and column b are independent of x_e . We consider two cases. If $i = r$ or $j = r$, then clearly $Q_W^{i,j} = Q_{W \setminus e}^{i,j}$. Otherwise,

$$Q_W^{i,j} = Q_{W \setminus e}^{i,j} + x_e \cdot Q_{W/e}^{i,j}.$$

This proves the claimed equalities also in this case (see Lemma 3.23) and the particular claim follows. \square

As an application of Lemma 3.23, we describe the behavior of configuration polynomials under 2-separations.

Proposition 3.27 (Configuration polynomials and 2-separations). *Let $W \subseteq \mathbb{K}^E$ be a realization of a connected matroid \mathbf{M} . Suppose that $E = E_1 \sqcup E_2$ is an (exact) 2-separation of \mathbf{M} . Then*

$$\psi_W = \psi_{W/E_1} \cdot \psi_{W|_{E_1}} + \psi_{W|_{E_2}} \cdot \psi_{W/E_2}.$$

Proof. We adopt the notation from [Tru92, §8.2]. Extend a basis $B_2 \in \mathcal{B}_{\mathbf{M}|_{E_2}}$ to a basis $B \in \mathcal{B}_{\mathbf{M}}$. Then W is the row span of a matrix (see [Tru92, (8.1.1)] and Remark 2.15)

$$A = \begin{pmatrix} I & 0 & A_1 & 0 \\ 0 & I & D & A_2' \end{pmatrix},$$

where the block columns are indexed by $B \setminus B_2, B_2, E_1 \setminus B, E_2 \setminus B_2$, and $\text{rk } D = 1$. After suitably ordering and scaling $B_2, E_1 \setminus B$ the lower rows of A , we may assume that

$$\begin{aligned} D &= (1 \ b)^t a_1, \\ a_1 &= (1 \ \cdots \ 1 \ 0 \ \cdots \ 0) \neq 0, \\ b &= (1 \ \cdots \ 1 \ 0 \ \cdots \ 0). \end{aligned}$$

The size of b and a_1 is determined by number of rows and columns of D , respectively. While b could be 0, at least one entry of a_1 is a 1. After suitable row operations and adjusting signs of B_2 , we can partition

$$(3.26) \quad A = \begin{pmatrix} I & 0 & 0 & A_1 & 0 \\ 0 & 1 & 0 & a_1 & a_2 \\ 0 & b^t & I & 0 & A_2 \end{pmatrix}.$$

Denote by $e \in E$ the index of the column $(0 \ 1 \ b)^t$. Let $X_1, x_e, X_2, X_1', X_2'$ be diagonal matrices of variables corresponding to the block columns of A . Then the configuration form of W becomes (see Remark 3.21)

$$Q_W = \begin{pmatrix} X_1 + A_1 X_1' A_1^t & A_1 X_1' a_1^t & 0 \\ a_1 X_1' A_1^t & x_e + a_1 X_1' a_1^t + a_2 X_2' a_2^t & x_e b + a_2 X_2' A_2^t \\ 0 & b^t x_e + A_2 X_2' a_2^t & b^t x_e b + X_2 + A_2 X_2' A_2^t \end{pmatrix},$$

which involves

$$\begin{aligned} Q_{W|_{E_1}} &= \begin{pmatrix} Q_{W/E_2} & A_1 X_1' a_1^t \\ a_1 X_1' A_1^t & a_1 X_1' a_1^t \end{pmatrix}, \\ Q_{W/E_2} &= X_1 + A_1 X_1' A_1^t, \\ Q_{W|_{E_2}} &= \begin{pmatrix} x_e + a_2 X_2' a_2^t & x_e b + a_2 X_2' A_2^t \\ b^t x_e + A_2 X_2' a_2^t & Q_{W/E_1} \end{pmatrix}, \\ Q_{W/E_1} &= b^t x_e b + X_2 + A_2 X_2' A_2^t. \end{aligned}$$

Laplace expansion of $\psi_W = \det Q_W$ (see Lemma 3.23) along the e th column yields the claimed formula. \square

Remark 3.28 (Configuration polynomials and handles). Let $W \subseteq \mathbb{K}^E$ be a realization of a connected matroid M , and let $H \in \mathcal{H}_M$ be a separating handle. By Lemma 2.4.(e), H is a 2-separation of M . Proposition 3.27 applied to $E = (E \setminus H) \sqcup H$ thus yields the statement of Corollary 3.13 in this case. \diamond

4. CONFIGURATION HYPERSURFACES

In this section, we establish our main results on Jacobian and second degeneracy schemes of realizations of connected matroids: the second degeneracy scheme is Cohen–Macaulay, the Jacobian scheme equidimensional, of codimension 3 (see Theorem 4.25). The second degeneracy scheme is reduced, the Jacobian scheme generically reduced if $\text{ch } \mathbb{K} \neq 2$ (see Theorem 4.25).

4.1. Commutative ring basics. In this subsection, we review the relevant preliminaries on equidimensionality and graded Cohen–Macaulayness using the books of Matsumura (see [Mat89]) and Bruns and Herzog (see [BH93]) as comprehensive references. For the benefit of the non-experts we provide detailed proofs. Further we relate generic reducedness for a ring and an associated graded ring (see Lemma 4.7).

4.1.1. Equidimensionality of rings. Let R be a Noetherian ring. We denote by $\text{Min Spec } R$ and $\text{Max Spec } R$ the sets of minimal and maximal elements of the set $\text{Spec } R$ of prime ideals of R with respect to inclusion. The subset $\text{Ass } R \subseteq \text{Spec } R$ of *associated primes* of R is finite and $\text{Min Spec } R \subseteq \text{Ass } R$ (see [Mat89, Thm. 6.5]).

One says that R is *catenary* if every saturated chain of prime ideals joining $\mathfrak{p}, \mathfrak{q} \in \text{Spec } R$ with $\mathfrak{p} \subseteq \mathfrak{q}$ has (maximal) length $\text{height}(\mathfrak{q}/\mathfrak{p})$ (see [Mat89, p. 31]). We say that R is *equidimensional* if it is catenary and $\forall \mathfrak{p} \in \text{Min Spec } R: \forall \mathfrak{m} \in \text{Max Spec } R: \mathfrak{p} \subseteq \mathfrak{m} \implies \text{height}(\mathfrak{m}/\mathfrak{p}) = \dim R$.

If R is a finitely generated \mathbb{K} -algebra, then these two conditions reduce to (see [BH93, Thm. 2.1.12] and [Mat89, Thm. 5.6])

$$\forall \mathfrak{p} \in \text{Min Spec } R: \dim(R/\mathfrak{p}) = \dim R.$$

We say that R is *pure-dimensional* if

$$\forall \mathfrak{p} \in \text{Ass } R: \dim(R/\mathfrak{p}) = \dim R,$$

which implies in particular that $\text{Ass } R = \text{Min Spec } R$. It follows that pure-dimensional finitely generated \mathbb{K} -algebras are equidimensional.

The following lemma applies to any equidimensional finitely generated \mathbb{K} -algebra.

Lemma 4.1 (Height bound for adding elements). *Let R be a Noetherian ring such that $R_{\mathfrak{m}}$ is equidimensional for all $\mathfrak{m} \in \text{Max Spec } R$.*

(a) *All saturated chains of primes in $\mathfrak{p} \in \text{Spec } R$ have length $\text{height } \mathfrak{p}$.*

(b) For any $\mathfrak{p} \in \text{Spec } R$, $x \in R$ and $\mathfrak{q} \in \text{Spec } R$ minimal over $\mathfrak{p} + \langle x \rangle$,
 $\text{height } \mathfrak{q} \leq \text{height } \mathfrak{p} + 1$.

Proof.

(a) Take two such chains of length n and n' starting at minimal primes \mathfrak{p}_0 and \mathfrak{p}'_0 , respectively. Extend both by a saturated chain of primes of length m containing \mathfrak{p} and ending in a maximal ideal \mathfrak{m} . Since $R_{\mathfrak{m}}$ is equidimensional by hypothesis, these extended chains have length $n + m = n' + m$. Therefore, the two chains have length $n = n'$.

(b) By Krull's principal ideal theorem, $\text{height}(\mathfrak{q}/\mathfrak{p}) \leq 1$. Take a chain of primes in \mathfrak{p} of length $\text{height } \mathfrak{p}$ and extend it by \mathfrak{q} if $\mathfrak{p} \neq \mathfrak{q}$. By (a), this extended chain has length $\text{height } \mathfrak{q}$ and the claim follows. \square

Lemma 4.2 (Equidimensional finitely generated algebras and localization). *Let R be an equidimensional finitely generated \mathbb{K} -algebra and $x \in R$. If $R_x \neq 0$, then R_x is equidimensional of dimension $\dim R_x = \dim R$.*

Proof. Any minimal prime ideal of R_x is of the form \mathfrak{p}_x where $\mathfrak{p} \in \text{Min Spec } R$ with $x \notin \mathfrak{p}$. By the Hilbert Nullstellensatz (see [Mat89, Thm. 5.5]),

$$\bigcap \text{Max } V(\mathfrak{p}) = \mathfrak{p}.$$

This yields an $\mathfrak{m} \in \text{Max Spec } R$ such that $\mathfrak{p} \subseteq \mathfrak{m} \not\ni x$ and hence $\mathfrak{p}_x \subseteq \mathfrak{m}_x \in \text{Max Spec } R_x$. Since R and hence R_x is a finitely generated \mathbb{K} -algebra,

$$\dim(R_x/\mathfrak{p}_x) = \text{height}(\mathfrak{m}_x/\mathfrak{p}_x) = \text{height}(\mathfrak{m}/\mathfrak{p}) = \dim R$$

by equidimensionality of R . The claim follows. \square

4.1.2. *Generic reducedness.* The following types of Artinian local rings coincide: field, regular ring, integral domain and reduced ring (see [Mat89, Thms. 2.2, 14.3]). A Noetherian ring R is *generically reduced* if the Artinian local ring $R_{\mathfrak{p}}$ is reduced for all $\mathfrak{p} \in \text{Min Spec } R$ (see [Mat89, Exc. 5.2]). This is equivalent to R satisfying Serre's condition (R_0) . We use the same notions for the associated affine scheme $\text{Spec } R$.

Definition 4.3 (Generic reducedness). We call a Noetherian scheme X *generically reduced along a subscheme Y* if X is reduced at all generic points specializing to a point of Y . If $X = \text{Spec } R$ is an affine scheme, then we use the same notions for the Noetherian ring R .

Lemma 4.4 (Reducedness and purity). *A Noetherian ring R is reduced if it is generically reduced and pure-dimensional.*

Proof. Since R is pure-dimensional, $\text{Ass } R = \text{Min Spec } R$, and hence, R becomes a subring of localizations (see [Mat89, Thm. 6.1.(i)])

$$R \hookrightarrow \bigoplus_{\mathfrak{p} \in \text{Ass } R} R_{\mathfrak{p}} = \bigoplus_{\text{Min Spec } R} R_{\mathfrak{p}}.$$

The latter ring is reduced since R is generically reduced, and the claim follows. \square

Lemma 4.5 (Reducedness and reduction). *Let (R, \mathfrak{m}) be a local Noetherian ring. Suppose that R/tR is reduced for a system of parameters t . Then R is regular and, in particular, an integral domain and reduced.*

Proof. By hypothesis, R/tR is local Artinian with maximal ideal \mathfrak{m}/tR . Reducedness makes R/tR a field, and hence, $\mathfrak{m} = tR$. By definition, this means that R is regular. In particular, R is an integral domain and reduced (see [Mat89, Thm. 14.3]). \square

Definition 4.6 (Rees algebras). Let R be a ring and $I \trianglelefteq R$ an ideal. The (extended) Rees algebra is the $R[t]$ -algebra (see [HS06, Def. 5.1.1])

$$\text{Rees}_I R := R[t, It^{-1}] \subseteq R[t^{\pm 1}].$$

The associated graded algebra is the R/I -algebra

$$\text{gr}_I R := \bigoplus_{i=0}^{\infty} I^i/I^{i+1}.$$

Lemma 4.7 (Generic reducedness from associated graded ring). *Let R be a Noetherian d -dimensional ring, $I \trianglelefteq R$ an ideal, $S := \text{Rees}_I R$ and $\bar{R} := \text{gr}_I R$.*

- (a) *Suppose R is an equidimensional finitely generated \mathbb{K} -algebra. Then S is a $(d+1)$ -equidimensional finitely generated \mathbb{K} -algebra.*
- (b) *If S is $(d+1)$ -equidimensional and $I \neq R$, then \bar{R} is d -equidimensional.*
- (c) *If S is equidimensional and \bar{R} is generically reduced, then R is generically reduced along $V(I)$.*

Proof. There are ring homomorphisms

$$R \rightarrow R[t] \rightarrow S \rightarrow S/tS \cong \bar{R}.$$

Since R is Noetherian, I is finitely generated and S finite type over R .

(a) If R is an integral domain, then so are $S \subseteq R[t^{\pm 1}]$. By definition, formation of the Rees ring commutes with base change. After base change to R/\mathfrak{p} for some $\mathfrak{p} \in \text{Min Spec } R$, we may assume that R is a d -dimensional integral domain. Then S is a $(d+1)$ -dimensional integral domain (see [HS06, Thm. 5.1.4]). Since S is a finitely generated \mathbb{K} -algebra (as R is one), S is equidimensional.

(b) Multiplication by t is injective on $R[t^{\pm 1}]$ and hence on S . If $I \neq R$, then $S/tS \cong \bar{R} \neq 0$ and t is an S -sequence. Since S is $(d+1)$ -equidimensional, \bar{R} is d -equidimensional by Krull's principal ideal theorem.

(c) Let $\mathfrak{p} \in \text{Min Spec } R$ and consider the extension $\mathfrak{p}[t^{\pm 1}] \in \text{Spec } R[t^{\pm 1}]$. Then (see [HS06, p. 96])

$$t \notin \tilde{\mathfrak{p}} := \mathfrak{p}[t^{\pm 1}] \cap S \in \text{Min Spec } S$$

and hence

$$(4.1) \quad S_{\tilde{\mathfrak{p}}} = (S_t)_{\tilde{\mathfrak{p}}_t} = R[t^{\pm 1}]_{\mathfrak{p}[t^{\pm 1}]}.$$

Since $\mathfrak{p}[t^{\pm 1}] \cap R = \mathfrak{p}$, the map $R \rightarrow R[t^{\pm 1}]$ induces an injection

$$(4.2) \quad R_{\mathfrak{p}} \hookrightarrow R[t^{\pm 1}]_{\mathfrak{p}[t^{\pm 1}]}.$$

To check injectivity, consider $R_{\mathfrak{p}} \ni x/1 \mapsto 0 \in R[t^{\pm 1}]_{\mathfrak{p}[t^{\pm 1}]}$. Then $0 = xy \in R[t^{\pm 1}]$ for some $y = \sum_i y_i t^i \in R[t^{\pm 1}] \setminus \mathfrak{p}[t^{\pm 1}]$. Then $0 = xy_i \in R$ for all i and $y_j \in R \setminus \mathfrak{p}$ for some j . It follows that $0 = x/1 \in R_{\mathfrak{p}}$. Combining (4.1) and (4.2) reducedness of $R_{\mathfrak{p}}$ follows from reducedness of $S_{\tilde{\mathfrak{p}}}$.

Suppose now that $V(\mathfrak{p}) \cap V(I) \neq \emptyset$ and hence (the subscript denoting graded parts)

$$R \neq \mathfrak{p} + I = \tilde{\mathfrak{p}}_0 + (tS)_0 = (\tilde{\mathfrak{p}} + tS)_0$$

implies that $\tilde{\mathfrak{p}} + tS \neq S$. Let $\mathfrak{q} \in \text{Spec } S$ be a minimal prime ideal over $\tilde{\mathfrak{p}} + tS$. No minimal prime ideal of S contains the S -sequence $t \in \mathfrak{q}$. By Lemma 4.1.(b), height $\mathfrak{q} = 1$ and \mathfrak{q} is minimal over t . This makes t a parameter of the localization $S_{\mathfrak{q}}$. Under $S/tS \cong \bar{R}$, the minimal prime ideal $\mathfrak{q}/tS \in \text{Spec}(S/tS)$ corresponds to a minimal prime ideal $\bar{\mathfrak{q}} \in \text{Spec } \bar{R}$. Suppose that \bar{R} is generically reduced. Then

$$S_{\mathfrak{q}}/tS_{\mathfrak{q}} = (S/tS)_{\mathfrak{q}/tS} \cong \bar{R}_{\bar{\mathfrak{q}}}$$

is reduced. By Lemma 4.5, $S_{\mathfrak{q}}$ and hence its localization $(S_{\mathfrak{q}})_{\tilde{\mathfrak{p}}_{\mathfrak{q}}} = S_{\tilde{\mathfrak{p}}}$ is reduced. Then also $R_{\mathfrak{p}}$ is reduced, as shown before. \square

4.1.3. *Graded Cohen–Macaulay rings.* Let (R, \mathfrak{m}) be a Noetherian $*$ local ring (see [BH93, Def. 1.5.13]). By definition, this means that R is a graded ring with unique maximal graded ideal \mathfrak{m} . For any $\mathfrak{p} \in \text{Spec } R$, denote by $\mathfrak{p}^* \in \text{Spec } R$ the maximal graded ideal contained in \mathfrak{p} (see [BH93, Lem. 1.5.6.(a)]). For any $\mathfrak{p} \in \text{Spec } R$, there is a chain of maximal length of graded prime ideals strictly contained in \mathfrak{p} (see [BH93, Lem. 1.5.8]). If $\mathfrak{m} \notin \text{Max Spec } R$, then such a chain for $\mathfrak{n} \in \text{Max Spec } R$ ends with $\mathfrak{m} \subsetneq \mathfrak{n}$. It follows that

$$(4.3) \quad \dim R = \begin{cases} \dim R_{\mathfrak{m}} & \text{if } \mathfrak{m} \in \text{Max Spec } R, \\ \dim R_{\mathfrak{m}} + 1 & \text{if } \mathfrak{m} \notin \text{Max Spec } R. \end{cases}$$

For any proper graded ideal $I \triangleleft R$ also $(R/I, \mathfrak{m}/I)$ is $*$ local and

$$(4.4) \quad \mathfrak{m} \in \text{Max Spec } R \iff \mathfrak{m}/I \in \text{Max Spec}(R/I).$$

Any associated prime $\mathfrak{p} \in \text{Ass } R$ is graded (see [BH93, Lem. 1.5.6.(b).(ii)]) and hence $\mathfrak{p} \subseteq \mathfrak{m}$. This yields a bijection (see [Mat89, Thm. 6.2])

$$(4.5) \quad \text{Ass } R \rightarrow \text{Ass } R_{\mathfrak{m}}, \quad \mathfrak{p} \mapsto \mathfrak{p}_{\mathfrak{m}}.$$

If $I \trianglelefteq R$ is a graded ideal and $\mathfrak{p} \in \text{Spec } R$ minimal over I , then $\mathfrak{p}/I \in \text{Min Spec}(R/I) \subseteq \text{Ass}(R/I)$, and hence, \mathfrak{p} is graded.

The following lemma shows in particular that $*$ local Cohen–Macaulay rings are pure- and equidimensional.

Lemma 4.8 (Height and codimension). *Let (R, \mathfrak{m}) be a *local Cohen–Macaulay ring and $I \trianglelefteq R$ a graded ideal. Then R is pure-dimensional and*

$$(4.6) \quad \text{height } I = \text{codim } I.$$

In particular, R/I is equidimensional if and only if $\text{height } \mathfrak{p} = \text{codim } I$ for all minimal $\mathfrak{p} \in \text{Spec } R$ over I .

Proof. The *local ring (R, \mathfrak{m}) is Cohen–Macaulay if and only if the localization $R_{\mathfrak{m}}$ is Cohen–Macaulay (see [BH93, Exc. 2.1.27.(c)]). In particular, $R_{\mathfrak{m}}$ is pure-dimensional (see [BH93, Prop. 1.2.13]) and (see [BH93, Cor. 2.1.4])

$$(4.7) \quad \text{height } I_{\mathfrak{m}} = \text{codim } I_{\mathfrak{m}}$$

Using (4.3), (4.4) for $I = \mathfrak{p}$ and bijection (4.5), it follows that R is pure-dimensional:

$$\begin{aligned} \forall \mathfrak{p} \in \text{Ass } R: \dim R &= \begin{cases} \dim R_{\mathfrak{m}} & \text{if } \mathfrak{m} \in \text{Max Spec } R, \\ \dim R_{\mathfrak{m}} + 1 & \text{if } \mathfrak{m} \notin \text{Max Spec } R, \end{cases} \\ &= \begin{cases} \dim(R_{\mathfrak{m}}/\mathfrak{p}_{\mathfrak{m}}) & \text{if } \mathfrak{m} \in \text{Max Spec } R, \\ \dim(R_{\mathfrak{m}}/\mathfrak{p}_{\mathfrak{m}}) + 1 & \text{if } \mathfrak{m} \notin \text{Max Spec } R, \end{cases} \\ &= \begin{cases} \dim(R/\mathfrak{p})_{\mathfrak{m}/\mathfrak{p}} & \text{if } \mathfrak{m} \in \text{Max Spec } R, \\ \dim(R/\mathfrak{p})_{\mathfrak{m}/\mathfrak{p}} + 1 & \text{if } \mathfrak{m} \notin \text{Max Spec } R, \end{cases} \\ &= \dim(R/\mathfrak{p}). \end{aligned}$$

Using (4.3) and (4.4), (4.6) follows from (4.7):

$$\begin{aligned} \text{height } I &= \text{height } I_{\mathfrak{m}} = \text{codim } I_{\mathfrak{m}} \\ &= \dim R_{\mathfrak{m}} - \dim(R_{\mathfrak{m}}/I_{\mathfrak{m}}) \\ &= \dim R_{\mathfrak{m}} - \dim(R/I)_{\mathfrak{m}/I} \\ &= \dim R - \dim(R/I) = \text{codim } I. \end{aligned}$$

Since R is Cohen–Macaulay, it is (universally) catenary (see [BH93, Thm. 2.1.12]). By (4.4) and the preceding discussion of chains of prime ideals in R/I and R/\mathfrak{p} , I is equidimensional if and only if $\dim(R/I) = \dim(R/\mathfrak{p})$ for all prime ideals $\mathfrak{p} \in \text{Spec } R$ minimal over I . The particular claim then follows by (4.6) for I and \mathfrak{p} . \square

4.2. Jacobian and degeneracy schemes. In this subsection, we associate Jacobian and second degeneracy schemes to a configuration. By results of Patterson and Kutz, their supports coincide and their codimension is at most 3.

For a Noetherian ring R , we consider the associated affine (Noetherian) *scheme* $\text{Spec } R$, whose underlying set consists of all prime ideals of R . We refer to elements of $\text{Min Spec } R$ as *generic points*, of $\text{Ass } R$

as *associated points*, and of $\text{Ass } R \setminus \text{Min Spec } R$ as *embedded points* of $\text{Spec } R$. An ideal $I \trianglelefteq R$ defines a *subscheme* $\text{Spec}(R/I) \subseteq \text{Spec } R$.

By abuse of notation we identify

$$\mathbb{K}^E = \text{Spec } \mathbb{K}[x].$$

Due to Lemma 4.8,

$$\text{codim}_{\mathbb{K}^E} \text{Spec}(\mathbb{K}[x]/I) = \text{height } I$$

for any graded ideal $I \trianglelefteq \mathbb{K}[x]$.

Definition 4.9 (Configuration schemes). Let $W \subseteq \mathbb{K}^E$ be a configuration. Then the subscheme

$$X_W := \text{Spec}(\mathbb{K}[x]/\langle \psi_W \rangle) \subseteq \mathbb{K}^E$$

is called the *configuration hypersurface* of W . In particular, $X_G := X_{W_G}$ is the *graph hypersurface* of G (see Definition 2.23). The ideal

$$J_W := \langle \psi_W \rangle + \langle \partial_e \psi_W \mid e \in E \rangle \trianglelefteq \mathbb{K}[x]$$

is the *Jacobian ideal* of ψ_W . We call the subschemes (see Definition 3.20)

$$\Sigma_W := \text{Spec}(\mathbb{K}[x]/J_W) \subseteq \mathbb{K}^E, \quad \Delta_W := \text{Spec}(\mathbb{K}[x]/M_W) \subseteq \mathbb{K}^E,$$

the *Jacobian scheme* of X_W and the *second degeneracy scheme* of Q_W .

Remark 4.10 (Degeneracy and non-smooth loci). If $\text{ch } \mathbb{K} \nmid \text{rk } M = \text{deg } \psi$ (see Remark 3.5), then ψ_W is a redundant generator of J_W due to the Euler identity. By Lemma 3.23, X_W^{red} and Δ_W^{red} are the first and second degeneracy loci of Q_W (see Definition 3.20), whereas Σ_W^{red} is the *non-smooth locus* of X_W over \mathbb{K} (see [Mat89, Thm. 30.3.(1)]). If \mathbb{K} is perfect, then Σ_W^{red} is the *singular locus* of X_W (see [Mat89, §28, Lem. 1]). \diamond

Remark 4.11 (Loops and line factors). Let $W \subseteq \mathbb{K}^E$ be a realization of matroid M . Suppose that e is a loop in M , that is, $e^\vee|_W = 0$. Then ψ_W and Q_W are independent of x_e (see Remark 3.5 and Definition 3.20)

$$X_W = X_{W \setminus e} \times \mathbb{A}^1, \quad \Sigma_W = \Sigma_{W \setminus e} \times \mathbb{A}^1, \quad \Delta_W = \Delta_{W \setminus e} \times \mathbb{A}^1. \quad \diamond$$

Lemma 4.12 (Inclusions of schemes). *For any configuration $W \subseteq \mathbb{K}^E$, there are inclusions of schemes $\Delta_W \subseteq \Sigma_W \subseteq X_W \subseteq \mathbb{K}^E$.*

Proof. By definition, $\psi_W \in J_W$ and hence the second inclusion. By Lemma 3.23, $\psi_W = \det Q_W \in M_W$ and hence $\partial_e \psi_W \in M_W$ for all $e \in E$. Thus, $J_W \subseteq M_W$ and the first inclusion follows. \square

Remark 4.13 (Schemes for matroids of small rank). Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid M .

(a) If $\text{rk } M \leq 1$, then $\psi_W = 1$ (see Remark 3.5) or $\psi_W \neq 0$ is a \mathbb{K} -linear form. In both cases, $\Sigma_W = \emptyset = \Delta_W$. If $\text{rk } M \geq 2$, then $\langle x \rangle \in \Sigma_W \neq \emptyset \neq \Delta_W \ni \langle x \rangle$.

(b) If $\text{rk } \mathbf{M} = 2$, then Δ_W is a \mathbb{K} -linear subspace of \mathbb{K}^E and hence an integral scheme. If $\text{ch } \mathbb{K} \neq 2$, the same holds for Σ_W due to the Euler identity (see Remark 4.10). Otherwise, the non-redundant quadratic generator ψ_W of J_W can make Σ_W non-reduced (see Example 4.14). \diamond

Example 4.14 (Schemes for the triangle). Let \mathbf{M} be a matroid on $E \in \mathcal{C}_M$ with $|E| = 3$ and hence $\text{rk } \mathbf{M} = |E| - 1 = 2$. Up to scaling and ordering $E = \{e_1, e_2, e_3\}$, any realization $W \subseteq \mathbb{K}^E$ of \mathbf{M} has the basis

$$w^1 := e_1 + e_3, \quad w^2 := e_2 + e_3.$$

With respect to this basis, we compute

$$Q_W = \begin{pmatrix} x_1 + x_3 & x_3 \\ x_3 & x_2 + x_3 \end{pmatrix},$$

$$M_W = \langle x_1 + x_3, x_2 + x_3, x_3 \rangle = \langle x_1, x_2, x_3 \rangle.$$

It follows that Δ_W is a reduced point.

On the other hand,

$$\psi_W = \det Q_W = x_1x_2 + x_1x_3 + x_2x_3,$$

$$J_W = \langle \psi_W, x_1 + x_2, x_1 + x_3, x_2 + x_3 \rangle.$$

The matrix expressing the linear generators $x_1 + x_2, x_1 + x_3, x_2 + x_3$ in terms of the variables x_1, x_2, x_3 has determinant 2. If $\text{ch } \mathbb{K} \neq 2$, then $J_W = \langle x_1, x_2, x_3 \rangle$ and Σ_W is a reduced point. Otherwise,

$$J_W = \langle \psi_W, x_1 - x_3, x_2 - x_3 \rangle = \langle x_1 - x_3, x_2 - x_3, x_3^2 \rangle$$

and Σ_W is a non-reduced point. \diamond

Lemma 4.15. *Consider two sets of variables $x = x_1, \dots, x_n$ and $y = y_1, \dots, y_m$. Let $0 \neq f \in I \trianglelefteq \mathbb{K}[x]$ and $0 \neq g \in J \trianglelefteq \mathbb{K}[y]$. Then*

$$f \cdot J[x] + I[y] \cdot g = \langle f, g \rangle \cap I[y] \cap J[x] \trianglelefteq \mathbb{K}[x, y].$$

Proof. For the non-obvious inclusion, take $h = af + bg \in I[y] \cap J[x]$. Since $f \in I[y]$, $bg \in I[y]$ and similarly $af \in J[x]$. Since $f \neq 0$ and J are in different variables, it follows that $a \in J[x]$ and similarly $b \in I[y]$. \square

Theorem 4.16 (Decompositions of schemes). *Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid \mathbf{M} without loops. Suppose that $\mathbf{M} = \bigoplus_{i=1}^n \mathbf{M}_i$ decomposes into connected components \mathbf{M}_i on E_i . Let $W = \bigoplus_{i=1}^n W_i$ be the induced decomposition into $W_i \subseteq \mathbb{K}^{E_i}$ (see Lemma 2.19). Then X_W is the reduced union of integral schemes $X_{W_i} \times \mathbb{K}^{E \setminus E_i}$, and Σ_W is the union of $\Sigma_{W_i} \times \mathbb{K}^{E \setminus E_i}$ and integral schemes $X_{W_i} \times X_{W_j} \times \mathbb{K}^{E \setminus (E_i \cup E_j)}$ for $i \neq j$. The same holds for Σ replaced by Δ . In particular, X_W is generically smooth over \mathbb{K} .*

Proof. Proposition 3.8 yields the claim on X_W (see Remark 3.5). For the claims on Σ_W and Δ_W , we may assume that $n = 2$ with \mathbf{M}_1 possibly disconnected. The general case then follows by induction on n .

By Proposition 3.8 and Definition 3.20, $\psi_W = \psi_{W_1} \cdot \psi_{W_2}$ and $Q_W = Q_{W_1} \oplus Q_{W_2}$. Then Lemma 4.15 yields

$$\begin{aligned} J_W &= \psi_{W_1} \cdot J_{W_2}[x_{E_1}] + J_{W_1}[x_{E_2}] \cdot \psi_{W_2} \\ &= \langle \psi_{W_1}, \psi_{W_2} \rangle \cap J_{W_1}[x_{E_2}] \cap J_{W_2}[x_{E_1}], \end{aligned}$$

and hence,

$$\Sigma_W = (X_{W_1} \times X_{W_2}) \cup (\Sigma_{W_1} \times \mathbb{K}^{E_2}) \cup (\mathbb{K}^{E_1} \times \Sigma_{W_2}).$$

The same holds for J and Σ replaced by M and Δ , respectively.

Suppose now that M is connected. By Proposition 3.12, $\psi_W \nmid \partial_e \psi_W$ for any $e \in E$ and hence $\Sigma_W \subsetneq X_W$. The particular claim follows. \square

Patterson proved the following result (see [Pat10, Thm. 4.1]). While Patterson assumes $\text{ch } \mathbb{K} = 0$ and excludes the generator $\psi_W \in J_W$, his proof works in general (see Remark 4.10). We give an alternative proof using Dodgson identities.

Theorem 4.17 (Non-smooth loci and second degeneracy schemes). *Let $W \subseteq \mathbb{K}^E$ be a configuration. Then there is an equality of reduced loci*

$$\Sigma_W^{\text{red}} = \Delta_W^{\text{red}}.$$

In particular, Σ_W and Δ_W have the same generic points, that is,

$$\text{Min } \Sigma_W = \text{Min } \Delta_W.$$

Proof. Order $E = \{e_1, \dots, e_n\}$ and pick a basis $w = (w^1, \dots, w^r)$ of W . We may assume that its coefficients with respect to e_1, \dots, e_r form an identity matrix, that is, $w_{e_j}^i = \delta_{i,j}$ for $i, j \in \{1, \dots, r\}$. For $i, j \in \{1, \dots, r\}$ denote by $Q_W^{\{i,j\},\{i,j\}}$ the minor of Q_W obtained by deleting rows and columns i, j . Then there are Dodgson identities (see Remark 3.21, Lemma 3.23 and [BEK06, Lem. 8.2])

$$\begin{aligned} (Q_W^{i,j})^2 &= Q_W^{i,j} \cdot Q_W^{j,i} = Q_W^{i,i} \cdot Q_W^{j,j} - \det Q_W \cdot Q_W^{\{i,j\},\{i,j\}} \\ &= \partial_i \psi_W \cdot \partial_j \psi_W - \psi_W \cdot Q_W^{\{i,j\},\{i,j\}} \in J_W \end{aligned}$$

for $i, j \in \{1, \dots, r\}$. In particular, any prime ideal $\mathfrak{p} \in \text{Spec } \mathbb{K}[x]$ over J_W contains M_W and hence $\Sigma_W^{\text{red}} \subseteq \Delta_W^{\text{red}}$. The opposite inclusion is due to Lemma 4.12. \square

Corollary 4.18 (Cremona isomorphism). *Let $W \subseteq \mathbb{K}^E$ be a configuration. Then the Cremona isomorphism $\mathbb{T}^E \cong \mathbb{T}^{E^\vee}$ identifies*

$$\begin{aligned} X_W \cap \mathbb{T}^E &\cong X_{W^\perp} \cap \mathbb{T}^{E^\vee}, \\ \Sigma_W \cap \mathbb{T}^E &\cong \Sigma_{W^\perp} \cap \mathbb{T}^{E^\vee}, \\ \Delta_W \cap \mathbb{T}^E &\cong \Delta_{W^\perp} \cap \mathbb{T}^{E^\vee}. \end{aligned}$$

In particular, Σ_W , Δ_W , Σ_{W^\perp} and Δ_{W^\perp} have the same generic points in $\mathbb{T}^E \cong \mathbb{T}^{E^\vee}$.

Proof. Propositions 3.10 and 3.25 yield the statements for X_W and Δ_W . The statement for Σ_W follows using that ζ_E (see (3.21)) identifies $x_e \partial_e = -x_{e^\vee} \partial_{e^\vee}$ for $e \in E$. The particular claim follows with Theorem 4.17. \square

Proposition 4.19 (Codimension bound). *Let $W \subseteq \mathbb{K}^E$ be a configuration. Then the codimensions of Σ_W and Δ_W in \mathbb{K}^E are bounded by*

$$\text{codim}_{\mathbb{K}^E} \Sigma_W = \text{codim}_{\mathbb{K}^E} \Delta_W \leq 3.$$

In case of equality, Δ_W is Cohen–Macaulay (and hence pure-dimensional) and Σ_W is equidimensional.

Proof. The equality of codimensions follows from Theorem 4.17. The scheme Δ_W is defined by the ideal M_W of submaximal minors of the symmetric matrix Q_W with entries in the Cohen–Macaulay ring $\mathbb{K}[x]$ (see [BH93, p. 2.1.9]). In particular, $\text{codim}_{\mathbb{K}^E} \Sigma_W = \text{grade } M_W$ (see [BH93, 2.1.2.(b)]). Kutz proved the claimed inequality and that M_W is a perfect ideal in case of equality (see [Kut74, Thm. 1]). In the latter case, $\mathbb{K}[x]/M_W = \mathbb{K}[\Delta_W]$ is a Cohen–Macaulay ring (see [BH93, Thm. 2.1.5.(a)]) and hence pure-dimensional (see Lemma 4.8). Then Σ_W is equidimensional by Theorem 4.17. \square

4.3. Generic points and codimension. In this subsection, we show that the Jacobian and second degeneracy schemes reach the codimension bound of 3 in case of connected matroids. The statements on codimension and Cohen–Macaulayness in our main result follow. In the process, we obtain a description of the generic points in relation with any non-disconnective handle.

Lemma 4.20 (Primes over the Jacobian ideal and handles). *Let $W \subseteq \mathbb{K}^E$ be a realization of a connected matroid \mathbf{M} , and let $H \in \mathcal{H}_{\mathbf{M}}$ be a proper handle.*

- (a) *For any $h \in H$, $x^{H \setminus \{h\}} \cdot \psi_{W \setminus H} \in J_W$.*
- (b) *For any $e, f \in H$ with $e \neq f$, $x^{H \setminus \{e, f\}} \cdot \psi_{W \setminus H} \in J_W + \langle x_e, x_f \rangle$.*
- (c) *For any $d \in H$ and $e \in E \setminus H$, $x^{H \setminus \{d\}} \cdot \partial_e \psi_{W \setminus H} \in J_W + \langle x_d \rangle$.*
- (d) *If $\mathfrak{p} \in \text{Spec } \mathbb{K}[x]$ with $J_W \subseteq \mathfrak{p} \not\subseteq \psi_{W \setminus H}$, then $\langle x_e, x_f, x_g \rangle \subseteq \mathfrak{p}$ for some $e, f, g \in H$ with $e \neq f \neq g \neq e$.*

Proof. By Remark 3.4 and Corollary 3.13, we may assume that

$$\psi_W = \sum_{h \in H} x^{H \setminus \{h\}} \cdot \psi_{W \setminus H} + x^H \cdot \psi_{W/H}$$

has the form (3.14).

(a) Using that ψ_W is a linear combination of square-free monomials (see Definition 3.2),

$$x^{H \setminus \{h\}} \cdot \psi_{W \setminus H} = \psi_W|_{x_h=0} = \psi_W - x_h \cdot \partial_h \psi_W \in J_W.$$

(b) This follows from

$$\begin{aligned} J_W \ni \partial_e \psi_W &= \sum_{h \in H} x^{H \setminus \{e, h\}} \cdot \psi_{W \setminus H} + x^{H \setminus \{e\}} \cdot \psi_{W/H} \\ &\equiv x^{H \setminus \{e, f\}} \cdot \psi_{W \setminus H} \pmod{\langle x_e, x_f \rangle}. \end{aligned}$$

(c) This follows from

$$\begin{aligned} J_W \ni \partial_e \psi_W &= \sum_{h \in H} x^{H \setminus \{h\}} \cdot \partial_e \psi_{W \setminus H} + x^H \cdot \partial_e \psi_{W/H} \\ &\equiv x^{H \setminus \{d\}} \cdot \partial_e \psi_{W \setminus H} \pmod{\langle x_d \rangle}. \end{aligned}$$

(d) By (a), the hypotheses force $x^{H \setminus \{h\}} \in \mathfrak{p}$ for all $h \in H$ and hence $\langle x_e, x_f \rangle \subseteq \mathfrak{p}$ for some $e, f \in H$ with $e \neq f$. Then $x^{H \setminus \{e, f\}} \in \mathfrak{p}$ by (b) and the claim follows. \square

Remark 4.21 (Primes over the Jacobian ideal and 2-separations). Let $W \subseteq \mathbb{K}^E$ be a realization of a connected matroid \mathbf{M} . Suppose that $E = E_1 \sqcup E_2$ is an (exact) 2-separation of \mathbf{M} . For $\{i, j\} = \{1, 2\}$, note that

$$d_i := \deg \psi_{W|_{E_i}} = \deg \psi_{W/E_j} + 1$$

and hence by Proposition 3.27

$$\begin{aligned} J_W \ni \psi_W &= \psi_{W/E_i} \cdot \psi_{W|_{E_i}} + \psi_{W|_{E_j}} \cdot \psi_{W/E_j}, \\ J_W \ni \sum_{e \in E_i} x_e \partial_e \psi_W &= d_i \cdot \psi_{W/E_i} \cdot \psi_{W|_{E_i}} + (d_i - 1) \cdot \psi_{W|_{E_j}} \cdot \psi_{W/E_j}. \end{aligned}$$

Subtracting $d_i \cdot \psi_W$ from the latter yields $\psi_{W|_{E_j}} \cdot \psi_{W/E_j} \in J_W$, for $j = 1, 2$. It follows that, for every prime ideal $\mathfrak{p} \in \text{Spec } \mathbb{K}[x]$ over J_W and every 2-separation F of \mathbf{M} , we have $\psi_{W|_F} \in \mathfrak{p}$ or $\psi_{W/F} \in \mathfrak{p}$. \diamond

Lemma 4.22 (Inductive codimension bound). *Let $W \subseteq \mathbb{K}^E$ be a realization of a connected matroid \mathbf{M} , and let $H \in \mathcal{H}_{\mathbf{M}}$ be a proper non-disconnective handle. Suppose that $\text{codim}_{\mathbb{K}^E \setminus H} \Sigma_{W \setminus H} = 3$. Then Σ_W is equidimensional of codimension*

$$\text{codim}_{\mathbb{K}^E} \Sigma_W = 3$$

with generic points of the following types:

- (a) $\mathfrak{p} = \langle x_e, x_f, x_g \rangle =: \mathfrak{p}_{e, f, g}$ for some $e, f, g \in H$ with $e \neq f \neq g \neq e$,
- (b) $\mathfrak{p} = \langle \psi_{W \setminus H}, x_d, x_h \rangle =: \mathfrak{p}_{H, d, h}$ for some $d, h \in H$ with $d \neq h$,
- (c) $\psi_{W \setminus H}, \psi_{W/H} \in \mathfrak{p} \nmid x_h$ for all $h \in H$.

Proof. Since H is non-disconnective, $\psi_{W \setminus H} \in \mathbb{K}[x_{E \setminus H}]$ is irreducible by Proposition 3.8. Since $d, h \in H$ with $d \neq h$, $\mathfrak{p}_{H, d, h} \in \text{Spec } \mathbb{K}[x]$ with height $\mathfrak{p}_{H, d, h} = 3$. The same holds for $\mathfrak{p}_{e, f, g}$.

By Lemma 4.8 and the dimension hypothesis, $J_{W \setminus H} \trianglelefteq \mathbb{K}[x_{E \setminus H}]$ has height 3. Thus, for any $d \in H$,

$$(4.8) \quad \text{height}(\langle J_{W \setminus H}, x_d \rangle) = \text{height } J_{W \setminus H} + 1 = 4.$$

In particular, $\Sigma_{W \setminus H} \neq \emptyset$ and hence $\Sigma_W \neq \emptyset$ by Remark 4.13.(a).

Let $\mathfrak{p} \in \text{Spec } \mathbb{K}[x]$ be any minimal prime ideal over J_W . By Lemma 4.8 and Proposition 4.19, it suffices to show for the equidimensionality that $\text{height } \mathfrak{p} \geq 3$. This follows in particular if \mathfrak{p} contains a prime ideal of type $\mathfrak{p}_{e,f,g}$ or $\mathfrak{p}_{H,d,h}$. By Lemma 4.20.(d), the former is the case if $\psi_{W \setminus H} \notin \mathfrak{p}$. We may thus assume that $\psi_{W \setminus H} \in \mathfrak{p}$. By Lemma 4.20.(c),

$$(4.9) \quad x^{H \setminus \{d\}} \cdot \partial_e \psi_{W \setminus H} \in \mathfrak{p} + \langle x_d \rangle.$$

for any $d \in H$ and $e \in E \setminus H$.

First suppose that $x_d \in \mathfrak{p}$ for some $d \in H$. If $x^{H \setminus \{d\}} \in \mathfrak{p}$, then \mathfrak{p} contains a prime ideal of type $\mathfrak{p}_{H,d,h}$ for some $h \in H \setminus \{d\}$. Otherwise, $\langle J_{W \setminus H}, x_d \rangle \subseteq \mathfrak{p}$ by (4.9) and hence $\text{height } \mathfrak{p} \geq 4$ by (4.8) (see Remark 4.23).

Now suppose that $x_h \notin \mathfrak{p}$ for all $h \in H$ and hence $\psi_{W/H} \in \mathfrak{p}$ by (3.11) and (3.13) in Corollary 3.13. Let $\mathfrak{q} \in \text{Spec } \mathbb{K}[x]$ be any minimal prime ideal over $\mathfrak{p} + \langle x_d \rangle$. By (4.9), \mathfrak{q} contains one of the ideals

$$(4.10) \quad \langle \psi_{W \setminus H}, \psi_{W/H}, x_d, x_h \rangle = \mathfrak{p}_{H,d,h} + \langle \psi_{W/H} \rangle, \quad \langle J_{W \setminus H}, x_d \rangle,$$

for some $h \in H \setminus \{d\}$. By Lemma 2.4.(b) and (e) (see Remark 3.5),

$$\begin{aligned} \deg \psi_{W/H} &= \text{rk}(M/H) = \text{rk } M - |H| \\ &= \text{rk } M - \text{rk}(H) = \text{rk}(M \setminus H) - \lambda_M(H) < \deg \psi_{W \setminus H} \end{aligned}$$

and hence $\psi_{W \setminus H} \notin \mathfrak{p}_{H,d,h}$ and $\psi_{W/H} \notin \mathfrak{p}_{H,d,h}$. Thus, both ideals in (4.10) have height at least 4 (see (4.9)) and hence $\text{height } \mathfrak{q} \geq 4$. It follows that $\text{height}(\mathfrak{p} + \langle x_d \rangle) \geq 4$ and then $\text{height } \mathfrak{p} \geq 3$ by Lemma 4.1.(b). \square

Remark 4.23. The case where $\text{height } \mathfrak{p} \geq 4$ in the proof of Lemma 4.22 does finally not occur due to the Cohen–Macaulayness of Δ_W achieved by the argument (see Proposition 3.8). \diamond

Lemma 4.24 (Generic points for circuits). *Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid M on $E \in \mathcal{C}_M$ with $|E| - 1 = \text{rk } M \geq 2$. Then Σ_W^{red} is the union of all codimension-3 coordinate subspaces of \mathbb{K}^E .*

Proof. We apply the strategy of the proof of Lemma 4.22. By Remark 4.13.(4.13), the rank hypothesis implies that $\Sigma_W \neq \emptyset$. Let $\mathfrak{p} \in \text{Spec } \mathbb{K}[x]$ be any minimal prime ideal over J_W . If $\psi_{W \setminus H} \notin \mathfrak{p}$ for some $E \neq H \in \mathcal{H}_M$, then Lemma 4.20.(d) yields $e, f, g \in H$ with $e \neq f \neq g \neq e$ such that $\langle x_e, x_f, x_g \rangle \subseteq \mathfrak{p}$. Otherwise, \mathfrak{p} contains $x^{E \setminus H} = \psi_{W \setminus H} \in \mathfrak{p}$ for all $E \neq H \in \mathcal{H}_M$ and hence all x_e where $e \in E$. (This can only occur if $|E| = 3$.) By Lemma 4.8 and Proposition 4.19, it follows that $\mathfrak{p} = \langle x_e, x_f, x_g \rangle$. By symmetry, all such triples $e, f, g \in E$ occur (see Example 3.7). \square

Theorem 4.25 (Cohen–Macaulayness of degeneracy schemes). *Let $W \subseteq \mathbb{K}^E$ be a realization of a connected matroid M of rank $\text{rk } M \geq 2$. Then Δ_W is Cohen–Macaulay (and hence pure-dimensional) and Σ_W is equidimensional, both of codimension 3 in \mathbb{K}^E .*

Proof. By Proposition 4.19, it suffices to show that $\text{codim}_{\mathbb{K}^E} \Sigma_W = 3$. Lemma 2.13 yields a circuit $C \in \mathcal{C}_M$ of size $|C| \geq 3$ and $\text{codim}_{\mathbb{K}^C} \Sigma_{W|C} = 3$ by Lemma 4.24. Proposition 2.8 yields a handle decomposition of M of length k with $F_1 = C$. By Lemma 4.22 and induction on k , then also $\text{codim}_{\mathbb{K}^E} \Sigma_W = 3$. \square

Corollary 4.26 (Types of generic points). *Let $W \subseteq \mathbb{K}^E$ be a realization of a connected matroid M of rank $\text{rk } M \geq 2$, and let $H \in \mathcal{H}_M$ be a non-disconnective handle such that $\text{rk}(M \setminus H) \geq 2$. Then all generic points of Σ_W and Δ_W are of the types listed in Lemma 4.22 with respect to H .*

Proof. Applying Theorem 4.25 to the matroid $M \setminus H$ with realization $W \setminus H$, the claim follows from Lemma 4.22 and Theorem 4.17. \square

Corollary 4.27 (Generic points for 3-connected matroids). *Let $W \subseteq \mathbb{K}^E$ be a realization of a 3-connected matroid M with $|E| > 3$ if $\text{rk } M \geq 2$. Then all generic points of Σ_W and Δ_W lie in \mathbb{T}^E , that is,*

$$\text{Min } \Sigma_W = \text{Min } \Delta_W \subseteq \mathbb{T}^E.$$

Proof. The equality is due to Theorem 4.17. We may assume that $\Sigma_W \neq \emptyset$ and hence $\text{rk } M \geq 2$ by Remark 4.13.(a). Let $\mathfrak{p} \in \text{Min } \Sigma_W$ be a generic point of Σ_W . For any $e \in E$, consider the 1-handle $H := \{e\} \in \mathcal{H}_M$. By Proposition 2.5 and Lemma 2.4.(e), H is non-disconnective with $\text{rk}(M \setminus H) = \text{rk } M \geq 2$. Corollary 4.26 forces \mathfrak{p} to be of type (c) in Lemma 4.22. It follows that $\mathfrak{p} \in \bigcap_{e \in E} D(x_e) = \mathbb{T}^E$. \square

4.4. Reducedness of degeneracy schemes. In this subsection, we prove the reducedness statement in our main result as outlined in §1.4.

Lemma 4.28 (Generic reducedness for the prism). *Let $W \subseteq \mathbb{K}^E$ be any realization of the prism matroid (see Definition 2.1). Then $\Delta_W \cap \mathbb{T}^E$ is an integral scheme of codimension 3, defined by 3 linear binomials, each supported in a corresponding handle. If $\text{ch } \mathbb{K} \neq 2$, then also $\Sigma_W \cap \mathbb{T}^E = \Delta_W \cap \mathbb{T}^E$.*

Proof. By Remark 3.22, we may assume that W is the realization from Lemma 2.25. A corresponding matrix of Q_W is given in Example 3.24. Reducing its entries modulo $\mathfrak{p} := \langle x_1 + x_2, x_3 + x_4, x_5 + x_6 \rangle$ makes all its 3×3 -minors 0. Therefore, $J_W \subseteq M_W \subseteq \mathfrak{p}$ by Lemma 4.12. Using the minors

$$\begin{aligned} Q_W^{2,3} &= (x_1 + x_2) \cdot (-x_3x_5), \\ Q_W^{2,4} &= (x_1 + x_2) \cdot (-x_3) \cdot (x_5 + x_6), \\ Q_W^{3,4} &= (x_1 + x_2) \cdot (x_3 + x_4) \cdot x_5, \\ Q_W^{4,4} &= (x_1 + x_2) \cdot (x_3 + x_4) \cdot (x_5 + x_6), \end{aligned}$$

one computes that

$$-Q_W^{2,3} + Q_W^{2,4} - Q_W^{3,4} + Q_W^{4,4} = (x_1 + x_2) \cdot x_4 x_6.$$

By symmetry, it follows that $x_2 x_4 x_6 \cdot \mathfrak{p} \subseteq M_W$ and hence

$$\Delta_W \cap D(x_2 x_4 x_6) = V(\mathfrak{p}) \cap D(x_2 x_4 x_6).$$

Using ψ_W from Example 3.17, one computes that

$$\begin{aligned} (x_2 \cdot (x_2 \partial_2 - 1) + x_4 x_6 \cdot (\partial_3 + \partial_5) + (x_4 + x_6) \cdot (1 - x_4 \partial_4 - x_6 \partial_6)) \psi_W \\ = 2 \cdot (x_1 + x_2) \cdot x_4^2 x_6^2. \end{aligned}$$

By symmetry, it follows that $2 \cdot x_2^2 x_4^2 x_6^2 \cdot \mathfrak{p} \subseteq J_W$ and hence

$$\Sigma_W \cap D(x_2 x_4 x_6) = V(\mathfrak{p}) \cap D(x_2 x_4 x_6).$$

if $\text{ch } \mathbb{K} \neq 2$. □

More details on the prism matroid can be found in Example 5.1.

Lemma 4.29 (Reduction and deletion of non-(co)loops). *Let $e \in E$ be a non-(co)loop in a matroid \mathbf{M} . For any $I \trianglelefteq \mathbb{K}[x]$ set*

$$\bar{I} := (I + \langle x_e \rangle) / \langle x_e \rangle \trianglelefteq \mathbb{K}[x] / \langle x_e \rangle = \mathbb{K}[x_{E \setminus \{e\}}].$$

Then $J_{W \setminus e} \subseteq \bar{J}_W$ and $M_{W \setminus e} = \bar{M}_W$ for any realization $W \subseteq \mathbb{K}^E$ of \mathbf{M} .

Proof. This follows from Proposition 3.12 and Lemma 3.26. □

Lemma 4.30 (Generic reducedness and deletion of non-(co)loops). *Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid \mathbf{M} , and let $e \in E$ be a non-(co)loop. Then $\Sigma_{W \setminus e} = \emptyset$ implies $\Sigma_W = \emptyset$. Suppose that $\text{Min } \Sigma_W \subseteq D(x_e)$ and that Σ_W and $\Sigma_{W \setminus e}$ are equidimensional of the same codimension. If $\Sigma_{W \setminus e}$ is generically reduced, then Σ_W is generically reduced. In this case, each $\mathfrak{p} \in \text{Min } \Sigma_W$ defines a non-empty subset $\gamma(\mathfrak{p}) \subseteq \text{Min } \Sigma_{W \setminus e}$ such that*

$$(4.11) \quad V(\mathfrak{p}) \cap V(x_e) = \bigcup_{\mathfrak{q} \in \gamma(\mathfrak{p})} V(\mathfrak{q}),$$

$$(4.12) \quad \mathfrak{p} \neq \mathfrak{p}' \implies \gamma(\mathfrak{p}) \cap \gamma(\mathfrak{p}') = \emptyset.$$

In particular, $|\text{Min } \Sigma_W| \leq |\text{Min } \Sigma_{W \setminus e}|$. The same statements hold for Σ replaced by Δ .

Proof. The subscheme $\Sigma_W \cap V(x_e) \subseteq \mathbb{K}^{E \setminus \{e\}}$ is defined by the ideal \bar{J}_W (see Lemma 4.29). By Lemma 4.29 and since J_W is graded,

$$\begin{aligned} \Sigma_{W \setminus e} = \emptyset &\iff J_{W \setminus e} = \mathbb{K}[x_{E \setminus \{e\}}] \implies \bar{J}_W = \mathbb{K}[x] / \langle x_e \rangle \\ &\iff J_W + \langle x_e \rangle = \mathbb{K}[x] \iff J_W = \mathbb{K}[x] \iff \Sigma_W = \emptyset \end{aligned}$$

which is the first claim.

Let $\mathfrak{p} \in \text{Min } \Sigma_W$ be a generic point of Σ_W . Considered as an element of $\text{Spec } \mathbb{K}[x]$ it is minimal over J_W . Since J_W and hence \mathfrak{p} is graded,

$\mathfrak{p} + \langle x_e \rangle \neq \mathbb{K}[x]$. Let $\mathfrak{q} \in \text{Spec } \mathbb{K}[x]$ be minimal over $\mathfrak{p} + \langle x_e \rangle$. By Lemma 4.29,

$$(4.13) \quad J_{W \setminus e} \subseteq \bar{J}_W \subseteq \bar{\mathfrak{q}}.$$

Since $x_e \notin \mathfrak{p}$ by hypothesis, Lemma 4.1 shows that

$$\begin{aligned} \text{height } \mathfrak{q} &= \text{height } \mathfrak{p} + 1, \\ \text{height } \bar{\mathfrak{q}} &= \text{height } \mathfrak{q} - \text{height } \langle x_e \rangle = \text{height } \mathfrak{p}. \end{aligned}$$

By the dimension hypothesis, Lemma 4.8 and (4.13), it follows that $\bar{\mathfrak{q}}$ is minimal over both $J_{W \setminus e}$ and \bar{J}_W . The former means that $\bar{\mathfrak{q}} \in \text{Min } \Sigma_{W \setminus e}$. The set $\gamma(\mathfrak{p})$ of all such $\bar{\mathfrak{q}}$ is non-empty and satisfies condition (4.11).

Denote by $t \in \mathbb{K}[\Sigma_W]$ the image of x_e . Then $\mathfrak{q} \notin \text{Min } \mathbb{K}[\Sigma_W]$ by hypothesis and \mathfrak{q} is minimal over t since $\bar{\mathfrak{q}}$ is minimal over \bar{J}_W . This makes t a parameter of the localization

$$R := \mathbb{K}[\Sigma_W]_{\mathfrak{q}}.$$

The inclusion (4.13) gives rise to a surjection of local rings

$$(4.14) \quad \mathbb{K}[\Sigma_{W \setminus e}]_{\bar{\mathfrak{q}}} \twoheadrightarrow \mathbb{K}[\Sigma_W \cap V(x_e)]_{\bar{\mathfrak{q}}} = R/tR.$$

Suppose now that $\Sigma_{W \setminus e}$ is generically reduced. Then $\mathbb{K}[\Sigma_{W \setminus e}]_{\bar{\mathfrak{q}}}$ is a field which makes (4.14) an isomorphism. By Lemma 4.5, R is then an integral domain with unique minimal prime ideal $\mathfrak{p}_{\bar{\mathfrak{q}}}$. Thus, $\mathbb{K}[\Sigma_W]_{\mathfrak{p}} = R_{\mathfrak{p}_{\bar{\mathfrak{q}}}}$ is reduced and \mathfrak{p} is uniquely determined by $\bar{\mathfrak{q}}$. This uniqueness is condition (4.12). The particular claim follows immediately.

The preceding arguments remain valid if Σ and J are replaced by Δ and M , respectively: Lemma 4.29 applies in both cases. \square

Lemma 4.31 (Initial forms and contraction of non-(co)loops). *Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid \mathbf{M} . Suppose $E = F \sqcup G$ is partitioned in such a way that \mathbf{M}/G is obtained from \mathbf{M} by successively contracting non-(co)loops. For any ideal $J \trianglelefteq \mathbb{K}[x]_{x^G} = \mathbb{K}[x_F, x_G^{\pm 1}]$, denote by J^{inf} the ideal generated by the lowest x_F -degree parts of the elements of J . Then $J_{W/G}[x_G^{\pm 1}] \subseteq (J^{\text{inf}})_{x^G}$ and $M_{W/G}[x_G^{\pm 1}] \subseteq (M_W^{\text{inf}})_{x^G}$.*

Proof. We iterate Proposition 3.12 and Lemma 3.26, respectively, to pass from W to W/G by successively contracting non-(co)loops $e \in G$. This yields a basis of W extending a basis w^1, \dots, w^s of W/G such that

$$(4.15) \quad \begin{aligned} \psi_W &= x^G \cdot \psi_{W/G} + p, \\ \partial_f \psi_W &= x^G \cdot \partial_f \psi_{W/G} + \partial_f p, \\ Q_W^{i,j} &= x^G \cdot Q_{W/G}^{i,j} + q_{i,j}, \end{aligned}$$

for all $f \in F$ and $i, j \in \{1, \dots, s\}$, where $p, q_{i,j} \in \mathbb{K}[x]$ are polynomials with no term divisible by x^G . Since ψ_W and $Q_W^{i,j}$ are homogeneous linear combinations of square-free monomials (see Definition 3.2 and Lemma 3.26), $x^G \cdot \psi_{W/G}$, $x^G \cdot \partial_f \psi_{W/G}$ and $x^G \cdot Q_{W/G}^{i,j}$ are the respective lowest x_F -degree parts in (4.15). The claimed inclusions follow. \square

Lemma 4.32 (Generic reducedness and contraction of non-(co)loops). *Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid \mathbf{M} . Suppose $E = F \sqcup G$ is partitioned in such a way that \mathbf{M}/G is obtained from \mathbf{M} by successively contracting non-(co)loops. Then $\Sigma_{W/G} = \emptyset$ implies $\Sigma_W \cap D(x^G) \cap V(x_F) = \emptyset$. Suppose that $\Sigma_W \cap D(x^G)$ and $\Sigma_{W/G}$ are equidimensional of the same codimension. If $\Sigma_{W/G}$ is generically reduced, then $\Sigma_W \cap D(x^G)$ is generically reduced along $V(x_F)$. The same statements hold for Σ replaced by Δ .*

Proof. Consider the ideal

$$\begin{aligned} I &:= \langle x_F \rangle \trianglelefteq \mathbb{K}[\Sigma_W \cap D(x^G)] =: R \\ &= \mathbb{K}[\Sigma_W]_{x^G} = (\mathbb{K}[x]/J_W)_{x^G} = \mathbb{K}[x_F, x_G^{\pm 1}]/(J_W)_{x^G}, \end{aligned}$$

R being equidimensional by hypothesis. With notation from Lemma 4.31

$$\begin{aligned} \bar{R} &= \text{gr}_I R = \text{gr}_I((\mathbb{K}[x]/J_W)_{x^G}) \cong (\text{gr}_{\langle x_F \rangle}(\mathbb{K}[x]/J_W))_{x^G} \\ &\cong (\mathbb{K}[x]/J_W^{\text{inf}})_{x^G} = \mathbb{K}[x_F, x_G^{\pm 1}]/(J_W^{\text{inf}})_{x^G}. \end{aligned}$$

Lemma 4.31 then yields the first claim:

$$\begin{aligned} \Sigma_{W/G} = \emptyset &\iff J_{W/G} = \mathbb{K}[x_F] \iff J_{W/G}[x_G^{\pm 1}] = \mathbb{K}[x_F, x_G^{\pm 1}] \\ &\implies (J_W^{\text{inf}})_{x^G} = \mathbb{K}[x_F, x_G^{\pm 1}] \iff \bar{R} = 0 \iff I = R \\ &\iff \Sigma_W \cap D(x^G) \cap V(x_F) = \emptyset. \end{aligned}$$

The latter equality makes the second claim vacuous.

We may thus assume that $I \neq R$. Lemma 4.31 yields a surjection

$$\begin{aligned} \pi: \mathbb{K}[\Sigma_{W/G} \times \mathbb{T}^G] &= (\mathbb{K}[x_F]/J_{W/G})[x_G^{\pm 1}] \\ &= \mathbb{K}[x_F, x_G^{\pm 1}]/(J_{W/G}[x_G^{\pm 1}]) \twoheadrightarrow \bar{R}. \end{aligned}$$

By Lemmas 4.2 and 4.7 and the dimension hypothesis, source and target are equidimensional of the same dimension and hence π^{-1} induces

$$\text{Min Spec } \bar{R} \subseteq \text{Min}(\Sigma_{W/G} \times \mathbb{T}^G).$$

Suppose now that $\Sigma_{W/G}$ and hence $\Sigma_{W/G} \times \mathbb{T}^G$ is generically reduced. For any $\mathfrak{p} \in \text{Min Spec } \bar{R}$, this makes $\mathbb{K}[\Sigma_{W/G} \times \mathbb{T}^G]_{\mathfrak{p}}$ a field and due to

$$\pi_{\mathfrak{p}}: \mathbb{K}[\Sigma_{W/G} \times \mathbb{T}^G]_{\mathfrak{p}} \twoheadrightarrow \bar{R}_{\mathfrak{p}}$$

also $\bar{R}_{\mathfrak{p}}$ is a field. It follows that \bar{R} is generically reduced. By Lemma 4.7, R is then generically reduced along $V(I)$. This means that $\Sigma_W \cap D(x^G)$ is generically reduced along $V(x_F)$.

The preceding arguments remain valid if Σ and J are replaced by Δ and M , respectively: Lemma 4.31 applies in both cases. \square

Lemma 4.33 (Generic reducedness for circuits). *Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid \mathbf{M} on $E \in \mathcal{C}_{\mathbf{M}}$ of rank $\text{rk } \mathbf{M} = |E| - 1 \geq 2$. Then Δ_W is generically reduced. If $\text{ch } \mathbb{K} \neq 2$, then also Σ_W is generically reduced.*

Proof. We proceed by induction on $|E|$. The case $|E| = 3$ is covered by Example 4.14; here we use $\text{ch } \mathbb{K} \neq 2$.

Suppose now that $|E| > 3$. Let $\mathbf{p} \in \text{Min } \Sigma_W$ be a generic point of Σ_W . By Lemma 4.24, $\mathbf{p} = \langle x_e, x_f, x_g \rangle$ for some $e, f, g \in H$ with $e \neq f \neq g \neq e$. Pick $d \in E \setminus \{e, f, g\}$. Then $E \setminus \{d\} \in \mathcal{C}_{\mathbf{M}/d}$ and hence $\Sigma_{W/d}$ is generically reduced by induction. By Lemmas 4.2 and 4.32, $\Sigma_W \cap D(x_d)$ is then along $V(x_{E \setminus \{d\}})$. By choice of d , $\langle x_{E \setminus \{d\}} \rangle \in V(\mathbf{p}) \cap D(x_d)$. In other words, $\mathbf{p} \in \text{Min}(\Sigma_W \cap D(x_d))$ specializes to a point in $V(x_{E \setminus \{d\}}) \cap D(x_d)$. Thus, Σ_W is reduced at \mathbf{p} . It follows that Σ_W is generically reduced.

By Theorem 4.17, Δ_W has the same generic points as Σ_W . Therefore, the preceding arguments remain valid if Σ is replaced by Δ . \square

Lemma 4.34 (Generic reducedness and contraction of non-maximal handles). *Let $W \subseteq \mathbb{K}^E$ be a realization of a connected matroid \mathbf{M} of rank $\text{rk } \mathbf{M} \geq 2$. Assume that $|\text{Max } \mathcal{H}_{\mathbf{M}}| \geq 2$ and set*

$$\hbar := |E| - |\text{Max } \mathcal{H}_{\mathbf{M}}| \geq 0.$$

Suppose that $\Sigma_{W'}$ is generically reduced for every realization $W' \subseteq \mathbb{K}^{E'}$ of every connected matroid \mathbf{M}' of rank $\text{rk } \mathbf{M}' \geq 2$ with $|E'| < |E|$.

- (a) *If $\hbar > 3$, then Σ_W is generically reduced.*
- (b) *If $\hbar > 2$ and $e \in E$, then Σ_W is reduced at all $\mathbf{p} \in \text{Min } \Sigma_W \cap V(x_e)$. The same statements hold for Σ replaced by Δ .*

Proof. Let $\mathbf{p} \in \text{Spec } \mathbb{K}[x]$ with height $\mathbf{p} = 3$. Pick a subset $F \subseteq E$ such that $|F \cap H'| = 1$ for all $H' \in \text{Max } \mathcal{H}_{\mathbf{M}}$. If possible, pick $F \cap H' = \{e\}$ such that $x_e \in \mathbf{p}$. If $\hbar > 3$, then by Lemma 4.1.(b)

$$(4.16) \quad \text{height}(\mathbf{p} + \langle x_F \rangle) \leq 3 + |F| = 3 + |\text{Max } \mathcal{H}_{\mathbf{M}}| < |E| = \text{height} \langle x \rangle.$$

If $\hbar > 2$ and $\mathbf{p} \in V(x_e)$, then (4.16) holds with 3 replaced by 2. In either case pick $\mathbf{q} \in \text{Spec } \mathbb{K}[x]$ such that

$$(4.17) \quad \mathbf{p} + \langle x_F \rangle \subseteq \mathbf{q} \subsetneq \langle x \rangle.$$

Add to F all $f \in E$ with $x_f \in \mathbf{q}$. This does not affect (4.17). Then $x_g \notin \mathbf{q}$ and hence $x_g \notin \mathbf{p}$ for all $g \in G := E \setminus F \neq \emptyset$. In other words,

$$(4.18) \quad \mathbf{p} \in D(x^G), \quad \mathbf{q} \in V(\mathbf{p}) \cap D(x^G) \cap V(x_F) \neq \emptyset.$$

By the initial choice of F , $G \cap H' \subsetneq H'$ for each $H' \in \text{Max } \mathcal{H}_{\mathbf{M}}$. By Lemma 2.4.(d), successively contracting all elements of G does, up to bijection, not affect circuits and maximal handles. In particular, \mathbf{M}/G is a connected matroid on the set F , obtained from \mathbf{M} by successively contracting non-(co)loops.

Since $|F| \geq |\text{Max } \mathcal{H}_{\mathbf{M}}| \geq 2$, connectedness implies that $\text{rk}(\mathbf{M}/G) \geq 1$. If $\text{rk}(\mathbf{M}/G) = 1$, then $\Sigma_{W/G} = \emptyset$ by Remark 4.13.(a). Then $\Sigma_W \cap D(x^G) \cap V(x_F) = \emptyset$ by Lemma 4.32 and hence $\mathbf{p} \notin \Sigma_W$ by (4.18).

Suppose now that $\mathbf{p} \in \Sigma_W$ and hence $\text{rk}(\mathbf{M}/G) \geq 2$. Then $\Sigma_{W/G}$ is generically reduced by hypothesis, and $\mathbf{p} \in \Sigma_W \cap D(x^G)$ specializes to

a point in $V(x_F) \cap D(x^G)$ by (4.18). By Theorem 4.25 and Lemma 4.2, Σ_W , $\Sigma_W \cap D(x^G)$ and $\Sigma_{W/G}$ are equidimensional of codimension 3. By Lemma 4.8, height $\mathfrak{p} = 3$ means that $\mathfrak{p} \in \text{Min } \Sigma_W$. By Lemma 4.32, Σ_W is thus reduced at \mathfrak{p} . The claims follow.

The preceding arguments remain valid if Σ is replaced by Δ . \square

Lemma 4.35 (Reducedness for connected matroids). *Let $W \subseteq \mathbb{K}^E$ be a realization of a connected matroid \mathbf{M} of rank $\text{rk } \mathbf{M} \geq 2$. Then Δ_W is reduced. If $\text{ch } \mathbb{K} \neq 2$, then Σ_W is generically reduced.*

Proof. By Theorem 4.25, Δ_W is pure-dimensional. By Lemma 4.4, Δ_W is thus reduced if it is generically reduced. By Lemma 4.12 and Theorem 4.17, the first claim follows if Σ_W is generically reduced.

Assume that $\text{ch } \mathbb{K} \neq 2$. We proceed by induction on $|E|$. By Lemma 4.33, Σ_W is generically reduced if $E \in \mathcal{C}_{\mathbf{M}}$; the base case where $|E| = 3$ needs $\text{ch } \mathbb{K} \neq 2$. Otherwise, by Proposition 2.8, \mathbf{M} has a handle decomposition of length $k \geq 2$. By Proposition 2.12, \mathbf{M} has $k + 1$ (disjoint) non-disconnective handles $H = H_1, \dots, H_\ell \in \mathcal{H}_{\mathbf{M}}$ with

$$(4.19) \quad \ell \geq k + 1 \geq 3.$$

Note that $H_1, \dots, H_\ell \in \text{Max } \mathcal{H}_{\mathbf{M}} \cap \mathcal{I}_{\mathbf{M}}$ by Lemma 2.4.(c) and (b). In particular, $\text{rk}(\mathbf{M} \setminus H) \neq 0$.

Suppose first that $H = \{h\}$. Then $\text{rk}(\mathbf{M} \setminus h) \geq 2$ by Remark 4.13.(a) and Lemma 4.30, and $\text{Min } \Sigma_W \subseteq D(x_h)$ by Corollary 4.26. By Theorem 4.25, both Σ_W and $\Sigma_{W \setminus h}$ are equidimensional of codimension 3. Thus, Σ_W is generically reduced by Lemma 4.30 and the induction hypothesis.

Suppose now that $|H_i| \geq 2$ for all $i = 1, \dots, \ell$, and set (see Lemma 4.34)

$$m := |\text{Max } \mathcal{H}_{\mathbf{M}}|, \quad \hbar := |E| - m.$$

If $\hbar > 3$, then Σ_W is generically reduced by Lemma 4.34.(a) and the induction hypothesis. Otherwise,

$$2\ell + (m - \ell) \leq \sum_{i=1}^{\ell} |H_i| + (m - \ell) \leq |E| = \hbar + m \leq 3 + m$$

and hence $2\ell \leq \sum_{i=1}^{\ell} |H_i| \leq 3 + \ell$. Comparing with (4.19) yields $\ell = 3$, $k = 2$ and $|H_i| = 2$ for $i = 1, 2, 3$. By Lemma 2.10, $E = H_1 \sqcup H_2 \sqcup H_3$ is then the handle partition of \mathbf{M} . In particular, $\hbar = 6 - 3 = 3 > 2$. By Lemma 2.25, \mathbf{M} must be the prism matroid.

Let now $\mathfrak{p} \in \text{Min } \Sigma_W$ be a generic point of Σ_W , with \mathbf{M} the prism matroid. If $\mathfrak{p} \in \mathbb{T}^E$, then Σ_W is reduced at \mathfrak{p} by Lemma 4.28; here we use $\text{ch } \mathbb{K} \neq 2$ again. Otherwise, $\mathfrak{p} \in V(x_e)$ for some $e \in E$. Then Σ_W is reduced at \mathfrak{p} by Lemma 4.34.(b) and the induction hypothesis.

The preceding arguments remain valid for arbitrary $\text{ch } \mathbb{K}$ if Σ is replaced by Δ . \square

Theorem 4.36 (Reducedness). *Let $W \subseteq \mathbb{K}^E$ be a realization of a matroid M . Then*

$$\Delta_W = \Sigma_W^{\text{red}}$$

is reduced. If $\text{ch } \mathbb{K} \neq 2$, then Σ_W is generically reduced.

Proof. By Theorem 4.16 and Lemma 4.35 (see Remarks 4.11 and 4.13.(a)), Δ_W is reduced and Σ_W is generically reduced if $\text{ch } \mathbb{K} \neq 2$. The claimed equality is then due to Theorem 4.17. \square

4.5. Integrality of degeneracy schemes. In this subsection, we prove the following companion result to Proposition 3.8 as outlined in §1.4.

Theorem 4.37 (Integrality for 3-connected matroids). *Let $W \subseteq \mathbb{K}^E$ be a realization of a 3-connected matroid M of rank $\text{rk } M \geq 2$. Then Δ_W is integral and hence Σ_W is irreducible.*

Proof. The claim on Δ_W follows from Remark 4.13.(a) and Lemmas 4.38 and 4.43 and Corollary 4.41. Theorem 4.17 yields the claim on Σ_W . \square

In the following, we use notation from Example 2.26.

Lemma 4.38 (Reduction to wheels and whirls). *It suffices to verify Theorem 4.37 for $M \in \{W_n, W^n\}$ with $n \geq 3$.*

Proof. Let M and W be as in Theorem 4.37. By Remark 4.13.(b) and Theorem 4.17, the claim holds if $\text{rk } M = 2$. If $|E| \leq 4$, then $M = U_{2,n}$ where $n \in \{3, 4\}$ (see [Oxl11, Tab. 8.1]) and hence $\text{rk } M = 2$. We may thus assume that $\text{rk } M \geq 3$ and $|E| \geq 5$.

The 3-connectedness hypothesis on M holds equivalently for M^\perp (see 2.10). By Corollaries 4.18 and 4.27, the Cremona isomorphism thus identifies

$$(4.20) \quad \mathbb{T}^E \supseteq \text{Min } \Delta_W = \text{Min } \Delta_{W^\perp} \subseteq \mathbb{T}^{E^\vee}.$$

It follows that integrality is equivalent for Δ_W and Δ_{W^\perp} . In particular, we may also assume that $\text{rk } M^\perp \geq 3$.

We proceed by induction on $|E|$. Suppose that M is not a wheel or a whirl. Since $\text{rk } M \geq 3$, Tutte's wheels-and-whirls theorem (see [Oxl11, Thm. 8.8.4]) yields an $e \in E$ such that $M \setminus e$ or M/e is again 3-connected. In the latter case, we replace W by W^\perp and use (2.11). We may thus assume that $M \setminus e$ is 3-connected. Then $\Delta_{W \setminus e}$ is integral by induction hypothesis. Note that $\text{Min } \Delta_W \subseteq D(x_e)$ by (4.20). By Theorem 4.25, Δ_W and $\Delta_{W \setminus e}$ are equidimensional of codimension 3. By Remark 4.13.(a) and Lemma 4.30, $\Delta_W \neq \emptyset$ and $|\text{Min } \Delta_W| \leq |\text{Min } \Delta_{W \setminus e}| = 1$. It follows that Δ_W is integral. \square

Lemma 4.39 (Turning wheels). *Let $W \subseteq \mathbb{K}^E$ be the realization of W_n from Lemma 2.27. Then the cyclic group \mathbb{Z}_n acts on X_W , Σ_W and Δ_W by “turning the wheel”, induced by the generator $1 \in \mathbb{Z}_n$ mapping*

$$(4.21) \quad s_i \mapsto s_{i+1}, \quad r_i \mapsto r_{i+1}, \quad w^i \mapsto w^{i+1}.$$

Proof. By Lemma 2.27, W has a basis $w = (w_1, \dots, w_n)$ where $w^i = s_i + r_i - r_{i-1}$ for all $i \in \mathbb{Z}_n$. The assignment (4.21) stabilizes $W \subseteq \mathbb{K}^E$. The resulting \mathbb{Z}_n -action stabilizes ψ_W and Q_W , and hence J_W and M_W . As a consequence, it induces an action on X_W , Σ_W and Δ_W . \square

The graph hypersurface of the n -wheel was described by Bloch, Esnault and Kreimer (see [BEK06, (11.5)]). We show that it is also the unique configuration hypersurface of the n -whirl.

Proposition 4.40 (Schemes for wheels and whirls). *Let $W \subseteq \mathbb{K}^E$ be any realization of $M \in \{W_n, W^n\}$ where $E = S \sqcup R$. Then there are coordinates $z'_1, \dots, z'_n, y_1, \dots, y_n$ on \mathbb{K}^E such that*

$$\psi_W = \det Q_n, \quad M_W = I_{n-1}(Q_n),$$

where

$$Q_n := \begin{pmatrix} z'_1 & y_1 & 0 & \cdots & \cdots & 0 & y_n \\ y_1 & z'_2 & y_2 & 0 & \cdots & \cdots & 0 \\ 0 & y_2 & z'_3 & y_3 & 0 & \cdots & 0 \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & y_{n-3} & z'_{n-2} & y_{n-2} & 0 \\ 0 & \cdots & \cdots & 0 & y_{n-2} & z'_{n-1} & y_{n-1} \\ y_n & 0 & \cdots & \cdots & 0 & y_{n-1} & z'_n \end{pmatrix}.$$

In particular, X_W , Σ_W and Δ_W depend only on n up to isomorphism.

Proof. We may assume that W is the realization from Lemma 2.27. Denote the coordinates on $\mathbb{K}^E = \mathbb{K}^{S \sqcup R}$ by

$$(4.22) \quad z_1, \dots, z_n, y_1, \dots, y_n := s_1^\vee, \dots, s_n^\vee, r_1^\vee, \dots, r_n^\vee,$$

and consider the \mathbb{K} -linear automorphism defined by

$$z'_1 := z_1 + y_1 + t^2 \cdot y_n, \quad z'_i := z_i + y_i + y_{i-1}, \quad i = 2, \dots, n.$$

Then Q_W is represented by the matrix

$$\begin{pmatrix} z'_1 & -y_1 & 0 & \cdots & \cdots & 0 & -t \cdot y_n \\ -y_1 & z'_2 & -y_2 & 0 & \cdots & \cdots & 0 \\ 0 & -y_2 & z'_3 & -y_3 & 0 & \cdots & 0 \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & \cdots & 0 & -y_{n-3} & z'_{n-2} & -y_{n-2} & 0 \\ 0 & \cdots & \cdots & 0 & -y_{n-2} & z'_{n-1} & -y_{n-1} \\ -t \cdot y_n & 0 & \cdots & \cdots & 0 & -y_{n-1} & z'_n \end{pmatrix}.$$

Suitable scaling of y_1, \dots, y_n turns this matrix into Q_n . The particular claim follows with Lemma 3.23. \square

Corollary 4.41 (Small wheels and whirls). *Theorem 4.37 holds for the matroids $M = W_3$ and $M = W^n$ for $n \leq 4$.*

Proof. Let W be any realization of M . By Theorem 4.36, Δ_W is reduced and it suffices to check irreducibility, replacing \mathbb{K} by its algebraic closure. By Proposition 4.40, we may assume that $\Delta_W = V(I_{k+1}(Q_n))$ for $k = n - 2$.

Consider the morphism of algebraic varieties of matrices

$$Y := \mathbb{K}^{n \times k} \rightarrow \{A \in \mathbb{K}^{n \times n} \mid A = A^t, \text{rk } A \leq k\} =: Z, \quad B \mapsto BB^t.$$

Let $y_{i,j}$ and $z_{i,j}$ be the coordinates on Y and Z , respectively. Then Δ_W identifies with $V(z_{1,3}, z_{2,4}) \subseteq Z$ for $n = 4$ and with Z itself for $n \leq 3$. Both the preimage Y of Z and for $n = 4$ the preimage

$$V(y_{1,1}y_{1,3} + y_{1,2}y_{2,3}, y_{2,1}y_{1,4} + y_{2,2}y_{2,4})$$

of $V(z_{1,3}, z_{2,4})$ are irreducible. It thus suffices to show that Y surjects onto Z , which holds for all $k \leq n$.

Let $A \in Z$ and $I \subseteq \{1, \dots, n\}$ with $|I| = \text{rk } A = k$ and rows $i \in I$ of A linearly independent. Apply row operations C to make the rows $i \notin I$ of CA zero. Then CAC^t is nonzero only in rows and columns $i \in I$. Modifying C to include further row operations turns CAC^t into a diagonal matrix. As \mathbb{K} is algebraically closed, $CAC^t = D^2$ where D has exactly k nonzero diagonal entries. Then $A = BB^t$ where $B := C^{-1}D$, considered as an element of Y by dropping zero columns. \square

Lemma 4.42 (Operations on wheels and whirls). *Let $M \in \{W_n, W^n\}$.*

(a) *The bijection*

$$E = S \sqcup R \rightarrow E^\vee, \quad s_i \mapsto r_i^\vee, \quad r_i \mapsto s_i^\vee,$$

identifies $M = M^\perp$.

Suppose now that n is not minimal for M to be defined, that is, $n > 3$ if $M = W_n$ and $n > 2$ if $M = W^n$.

- (b) *The matroid $M \setminus s_n$ is connected of rank $\text{rk}(M \setminus s_n) \geq 2$. Its handle partition consists of non-disconnective handles, the 2-handle $\{r_{n-1}, r_n\}$ and 1-handles.*
- (c) *The matroid M/r_n is connected of rank $\text{rk}(M/r_n) \geq 2$. Its handle partition consists of non-disconnective 1-handles.*
- (d) *We can identify $W_n \setminus s_n / r_n = W_{n-1}$ and $W^n \setminus s_n / r_n = W^{n-1}$.*

Proof.

(a) The self-duality claim is obvious (see [Oxl11, Prop. 8.4.4]).

(b) This follows from the description of connectedness in terms of circuits (see (2.5) and Example 2.26).

(c) This follows from the description of connectedness in terms of circuits (see (2.7) and Example 2.26).

(d) The operation $M \mapsto M \setminus s_n / r_n$ deletes the triangle $\{s_{n-1}, r_{n-1}, s_n\}$ and maps the triangle $\{s_n, r_n, s_1\}$ to $\{s_{n-1}, r_{n-1}, s_1\}$ (see (2.5) and (2.7)). By duality, it acts on triads in the same way (see (a) and (2.11)). Moreover, $R \in \mathcal{C}_{M \setminus s_n / r_n}$ is equivalent to $R \in \mathcal{C}_M$ and hence

$M = W_n$ (see (2.5), (2.7) and Example 2.26). The claim then follows using the characterization of wheels and whirl in terms of triangles and triads (see Example 2.26). \square

Lemma 4.43 (Induction on wheels and whirls). *Theorem 4.37 for $M = W_n$ and $M = W^n$ follows from the cases $n = 3$ and $n \leq 4$, respectively.*

Proof. Suppose that n is not minimal for $M \in \{W_n, W^n\}$ to be defined. Let W' be any realization of M/r_n . Then $W' \setminus s_n$ is a realization of

$$M/r_n \setminus s_n = M \setminus s_n / r_n = M_{n-1}$$

by Lemma 4.42.(d). By induction hypothesis and Corollary 4.27, $\Delta_{W' \setminus s_n}$ is integral with generic point in $\mathbb{T}^{E \setminus \{s_n, r_n\}}$. By Lemma 4.42.(c) and Corollary 4.26, $\text{Min } \Delta_{W'} \subseteq \mathbb{T}^{E \setminus \{r_n\}} \subseteq D(s_n)$. By Lemma 4.42.(c) and Theorems 4.25, $\Delta_{W'}$ and $\Delta_{W' \setminus s_n}$ are equidimensional of codimension 3. By Remark 4.13.(a) and Lemma 4.30, $\Delta_{W'}$ is then integral.

Let W be any realization of M and use the coordinates from (4.22). By Lemma 4.42.(b) and Corollary 4.26, $\Delta_{W \setminus s_n}$ has at most one generic point \mathfrak{q}' in $V(y_{n-1}, y_n)$ while all the others lie in $\mathbb{T}^{E \setminus \{s_n\}}$. By Corollary 4.18, the Cremona isomorphism identifies the latter with generic points of $\Delta_{(W \setminus s_n)^\perp}$ in $\mathbb{T}^{E^\vee \setminus \{s_n^\vee\}}$. Use (2.11) and Lemma 4.42.(a) to identify

$$(M \setminus s_n)^\perp = M^\perp / s_n^\vee = M / r_n, \quad E^\vee \setminus \{s_n^\vee\} = E \setminus \{r_n\},$$

and consider $(W \setminus s_n)^\perp$ as a realization W' of M/r_n . By the above, $\Delta_{W'}$ is integral with generic point in $\mathbb{T}^{E \setminus \{r_n\}}$. Thus, $\Delta_{W \setminus s_n}$ has a unique generic point \mathfrak{q} in $\mathbb{T}^{E \setminus \{s_n\}}$. To summarize,

$$(4.23) \quad \text{Min } \Delta_{W \setminus s_n} = \{\mathfrak{q}, \mathfrak{q}'\}, \quad \mathfrak{q} \in \mathbb{T}^{E \setminus \{s_n\}}, \quad \mathfrak{q}' \in V(y_{n-1}, y_n).$$

By Lemma 4.42.(b) and Theorems 4.25 and 4.36, Δ_W and $\Delta_{W \setminus s_n}$ are equidimensional of codimension 3 and reduced. It suffices to show that Δ_W is irreducible. By way of contradiction, suppose that $\mathfrak{p} \neq \mathfrak{p}'$ for some $\mathfrak{p}, \mathfrak{p}' \in \text{Min } \Delta_W$. By Corollary 4.27, $\text{Min } \Delta_W \subseteq \mathbb{T}^E \subseteq D(s_n)$. By Lemma 4.30 and (4.23), it follows that

$$\Delta_W = \{\mathfrak{p}, \mathfrak{p}'\}.$$

By (4.11) in Lemma 4.30, we may assume that $\sqrt{\mathfrak{p}} = \mathfrak{q}$ and $\sqrt{\mathfrak{p}'} = \mathfrak{q}'$ where $\bar{I} := (I + \langle z_n \rangle) / \langle z_n \rangle$.

Consider first the case where $M = W_n$ with $n \geq 4$. By Remark 3.22, we may assume that W is the realization from Lemma 2.27. By Lemma 4.39, the cyclic group \mathbb{Z}_n acts on $\{\mathfrak{p}, \mathfrak{p}'\}$ by ‘‘turning the wheel’’. If it acts identically, then $\sqrt{\mathfrak{p}'} + \langle z_i \rangle \supseteq \langle y_{i-1}, y_i \rangle$ for all $i = 1, \dots, n$ and hence

$$\sqrt{\mathfrak{p}' + \langle z_1, \dots, z_n \rangle} = \langle z_1, \dots, z_n, y_1, \dots, y_n \rangle.$$

Then $\text{height}(\mathfrak{p}' + \langle z_1, \dots, z_n \rangle) = 2n$ which implies $\text{height } \mathfrak{p}' \geq n > 3$ by Lemma 4.1.(b), contradicting Theorem 4.25 (see Lemma 4.8).

Otherwise, the generator $1 \in \mathbb{Z}_n$ switches the assignment $\mathbf{p} \mapsto \mathbf{q}$ and $\mathbf{p} \mapsto \mathbf{q}'$ and $n = 2m$ must be even. Then $\sqrt{\mathbf{p} + \langle z_{2i} \rangle} \supseteq \langle y_{2i-1}, y_{2i} \rangle$ for all $i = 1, \dots, m$ and hence

$$\sqrt{\mathbf{p} + \langle z_2, z_4, z_6, \dots, z_n \rangle} \supseteq \langle z_2, z_4, z_6, \dots, z_n, y_1, \dots, y_n \rangle.$$

This leads to a contradiction as before.

Consider now the case where $\mathbf{M} = \mathbf{W}^n$ with $n \geq 5$. For $i = 1, \dots, n$, denote by \mathbf{q}_i and \mathbf{q}'_i the generic points of $\Delta_{W \setminus s_i}$ as in (4.23). By the pigeonhole principle, one of \mathbf{p} and \mathbf{p}' , say \mathbf{p} , is assigned to \mathbf{q}'_i for 3 spokes s_i . In particular, \mathbf{p} is assigned to \mathbf{q}'_i and \mathbf{q}'_j for two non-adjacent spokes s_i and s_j . Then

$$\sqrt{\mathbf{p} + \langle z_i, z_j \rangle} \supseteq \langle z_i, z_j, y_{i-1}, y_i, y_{j-1}, y_j \rangle.$$

This leads to a contradiction as before. The claim follows. \square

Theorem 4.37 proves the “only if” part of the following conjecture.

Conjecture 4.44 (Irreducibility and 3-connectedness). *Let \mathbf{M} be a matroid of rank $\text{rk } \mathbf{M} \geq 2$ on E . Then \mathbf{M} is 3-connected if and only if, for some/any realization $W \subseteq \mathbb{K}^E$ of \mathbf{M} , both Δ_W and Δ_{W^\perp} are integral.*

5. EXAMPLES

In this section, we illustrate our results with examples of prism, whirl and uniform matroids.

Example 5.1 (Prism matroid). Consider the prism matroid \mathbf{M} (see Definition 2.1) with its unique realization W (see Lemma 2.25). Then

$$\psi_W = x_1 x_2 (x_3 + x_4)(x_5 + x_6) + x_3 x_4 (x_1 + x_2)(x_5 + x_6) + x_5 x_6 (x_1 + x_2)(x_3 + x_4)$$

by Example 3.17. By Lemma 4.28, Δ_W has the unique generic point

$$\langle x_1 + x_2, x_3 + x_4, x_5 + x_6 \rangle$$

in \mathbb{T}^6 . By Corollary 4.26, there can be at most 3 more generic points symmetric to

$$\langle x_1, x_2, \psi_{W \setminus \{1,2\}} \rangle = \langle x_1, x_2, x_3 x_4 x_5 + x_3 x_4 x_6 + x_3 x_5 x_6 + x_4 x_5 x_6 \rangle.$$

Over $\mathbb{K} = \mathbb{F}_2$, their presence is confirmed by a computation in **Singular** (see [Dec+18]). It reveals a total of 7 embedded points in Σ_W . There is $\langle x_1, \dots, x_6 \rangle$, and 3 symmetric to each of

$$\langle x_3, x_4, x_5, x_6 \rangle \quad \text{and} \quad \langle x_1, x_2, x_3 + x_4, x_5 + x_6 \rangle.$$

Moreover, Σ_W is not reduced at any generic point. Since the above associated primes are geometrically prime, the conclusions remain valid over any field \mathbb{K} with $\text{ch } \mathbb{K} = 2$.

A **Singular** computation over \mathbb{Q} shows that Σ_W has exactly the above associated points for any field \mathbb{K} with $\text{ch } \mathbb{K} = 0$ or $\text{ch } \mathbb{K} \gg 0$. We expect that this holds in fact for $\text{ch } \mathbb{K} \neq 2$.


To verify at least the presence of these associated points in Σ_W for $\text{ch } \mathbb{K} \neq 2$, we claim that

$$\begin{aligned} \langle x_1, x_2, \psi_{W \setminus \{1,2\}} \rangle &= J_W : 2((x_3 + x_4)x_5^2 - (x_3 + x_4)x_6^2), \\ \langle x_3, x_4, x_5, x_6 \rangle &= J_W : 2(x_1 + x_2)^2 x_4 x_6, \\ \langle x_1, x_2, x_3 + x_4, x_5 + x_6 \rangle &= J_W : 2x_2(x_3 + x_4)x_6^2, \\ \langle x_1, \dots, x_6 \rangle &= J_W : 2(x_1 + x_2)(x_3 + x_4)x_6. \end{aligned}$$

The colon ideals on the right hand side can be read off from a suitable Gröbner basis (see [GP08, Lems. 1.8.3, 1.8.10 and 1.8.12]). Using **Singular** we compute such a Gröbner basis over \mathbb{Z} which confirms our claim. There are no odd prime numbers dividing its leading coefficients. It is therefore a Gröbner basis over any field \mathbb{K} with $\text{ch } \mathbb{K} \neq 2$ and the argument remains valid. \diamond

Example 5.2 (Whirl matroid). Consider the whirl matroid $\mathbf{M} := W^3$ (see Example 2.26). It is realized by 6 points in \mathbb{P}^2 with the collinearities shown in Figure 4. Since \mathbf{M} contracts to the uniform matroid $\mathbf{U}_{2,4}$,

FIGURE 4. Points in \mathbb{P}^2 defining the whirl matroid W^3 .



\mathbf{M} is not regular (see [Oxl11, Thm. 6.6.6]). The configuration polynomial reflects this fact. Using the realization W of \mathbf{M} from Lemma 2.27 with $t = -1$, $\{s_1, s_2, s_3\} = \{1, 2, 3\}$ and $\{r_1, r_2, r_3\} = \{4, 5, 6\}$, we find

$$\begin{aligned} \psi_W &= x_1x_2x_3 + x_1x_3x_4 + x_2x_3x_4 + x_1x_2x_5 + x_1x_3x_5 + x_1x_4x_5 \\ &\quad + x_2x_4x_5 + x_3x_4x_5 + x_1x_2x_6 + x_2x_3x_6 + x_1x_4x_6 + x_2x_4x_6 \\ &\quad + x_3x_4x_6 + x_1x_5x_6 + x_2x_5x_6 + x_3x_5x_6 + 4x_4x_5x_6. \end{aligned}$$

Replacing in ψ_W the coefficient 4 of $x_4x_5x_6$ by a 1 yields the matroid polynomial $\psi_{\mathbf{M}}$ (see Remark 3.6).

By Theorem 4.25, the configuration hypersurface X_W defined by ψ_W has 3-codimensional non-smooth locus Σ_W^{red} . Using **Singular** (see [Dec+18]) we compute a Gröbner basis over \mathbb{Z} of the ideal of partial derivatives of $\psi_{\mathbf{M}}$. The only prime numbers dividing leading coefficients are 2, 3 and 5. For $\text{ch } \mathbb{K} \neq 2, 3, 5$, it is therefore a Gröbner basis over \mathbb{K} . From its leading exponents we calculate that the non-smooth locus of the hypersurface defined by $\psi_{\mathbf{M}}$ has codimension 4 (see [GP08, Cor. 5.3.14]). By further **Singular** computations, this codimension is 4 for $\text{ch } \mathbb{K} = 2, 5$, and 3 for $\text{ch } \mathbb{K} = 3$. \diamond

Example 5.3 (Uniform rank-3 matroid). Suppose that $\text{ch } \mathbb{K} \neq 2, 3$. Then the configuration $W = \langle w^1, w^2, w^3 \rangle \subseteq \mathbb{K}^3$ defined by

$$(w_j^i)_{i,j} = \begin{pmatrix} 1 & 0 & 0 & 1 & 2 & 3 \\ 0 & 1 & 0 & 2 & 3 & 4 \\ 0 & 0 & 1 & 2 & 6 & 12 \end{pmatrix}$$

realizes the uniform matroid $U_{3,6}$ (see Example 2.20). The entries of $Q_w = (q_{i,j})_{i,j}$ satisfy the linear dependence relation (see Remark 3.21)

$$q_{1,2} + q_{1,3} = q_{2,3}.$$

By Lemma 3.23, ψ_W thus depends on fewer than 6 variables. More precisely, a **Singular** computation shows that Σ_W has Betti numbers $(1, 5, 10, 10, 5, 1)$, is not reduced and hence not Cohen–Macaulay.

Now, take W' to be a generic realization of $U_{3,6}$. Then the entries of $Q_{W'}$ with indices (i, j) where $i \leq j$ are linearly independent (see [BCK16, Prop. 6.4]), and $\Sigma_{W'}$ is reduced Cohen–Macaulay with Betti numbers $(1, 6, 8, 3)$. So basic geometric properties of the configuration hypersurface X_W are not determined by the matroid M , but depend on the realization W . \diamond

Example 5.4 (Uniform rank-2 matroid). Suppose that $\text{ch } \mathbb{K} \neq 2$ and consider the uniform matroid $U_{2,n}$ for $n \geq 3$ (see Examples 2.2 and 3.7.(c)). A realization W of $U_{2,n}$ is spanned by two vectors $w^1, w^2 \in \mathbb{K}^n$ for which (see Example 2.20)

$$c_{W,\{i,j\}} = \det \begin{pmatrix} w_i^1 & w_j^1 \\ w_i^2 & w_j^2 \end{pmatrix}^2 \neq 0,$$

for $1 \leq i < j \leq n$. Then

$$\psi_W = \sum_{1 \leq i < j \leq n} c_{W,\{i,j\}} \cdot x_i \cdot x_j,$$

and the ideal J_W is generated by n linear forms. These forms may be written as the rows of the Hessian matrix

$$H_W := H_{\psi_W} = (c_{W,\{i,j\}})_{i,j},$$

where by convention $c_{W,\{i,i\}} = 0$. Since uniform matroids are connected, Theorem 4.25 implies that H_W has rank exactly 3.

For $n \geq 4$, this amounts to a classical-looking linear algebra fact: suppose that $A = (a_{i,j}^2)_{i,j} \in \mathbb{K}^{n \times n}$ is a matrix with squared entries. Then its 4×4 minors are zero provided that the numbers $a_{i,j}$ satisfy the Plücker relations defining the Grassmannian $\text{Gr}_{2,n}$. An elementary direct proof was shown to us by Darij Grinberg (see [Gri18]). \diamond

REFERENCES

- [AM11a] Paolo Aluffi and Matilde Marcolli. “Algebraic-geometric Feynman rules”. In: *Int. J. Geom. Methods Mod. Phys.* 8.1 (2011), pp. 203–237.

- [AM11b] Paolo Aluffi and Matilde Marcolli. “Feynman motives and deletion-contraction relations”. In: *Topology of algebraic varieties and singularities*. Vol. 538. Contemp. Math. Amer. Math. Soc., Providence, RI, 2011, pp. 21–64.
- [AOV18] Nima Anari, Shayan Oveis Gharan, and Cynthia Vinzant. “Log-concave polynomials, entropy, and a deterministic approximation algorithm for counting bases of matroids”. In: *59th Annual IEEE Symposium on Foundations of Computer Science—FOCS 2018*. IEEE Computer Soc., Los Alamitos, CA, 2018, pp. 35–46.
- [BB03] Prakash Belkale and Patrick Brosnan. “Matroids, motives, and a conjecture of Kontsevich”. In: *Duke Math. J.* 116.1 (2003), pp. 147–188.
- [BCK16] Cristiano Bocci, Enrico Carlini, and Joe Kileel. “Hadamard products of linear spaces”. In: *J. Algebra* 448 (2016), pp. 595–617.
- [BEK06] Spencer Bloch, Hélène Esnault, and Dirk Kreimer. “On motives associated to graph polynomials”. In: *Comm. Math. Phys.* 267.1 (2006), pp. 181–225.
- [BH93] Winfried Bruns and Jürgen Herzog. *Cohen-Macaulay rings*. Vol. 39. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1993, pp. xii+403.
- [BK97] D. J. Broadhurst and D. Kreimer. “Association of multiple zeta values with positive knots via Feynman diagrams up to 9 loops”. In: *Phys. Lett. B* 393.3-4 (1997), pp. 403–412.
- [Bro15] Francis Brown. *Periods and Feynman amplitudes*. 2015. arXiv: [1512.09265](https://arxiv.org/abs/1512.09265).
- [Bro17] Francis Brown. “Feynman amplitudes, coaction principle, and cosmic Galois group”. In: *Commun. Number Theory Phys.* 11.3 (2017), pp. 453–556.
- [BS12] Francis Brown and Oliver Schnetz. “A K3 in ϕ^4 ”. In: *Duke Math. J.* 161.10 (2012), pp. 1817–1862.
- [BW10] Christian Bogner and Stefan Weinzierl. “Feynman graph polynomials”. In: *Internat. J. Modern Phys. A* 25.13 (2010), pp. 2585–2618.
- [CH96] Collette R. Coullard and Lisa Hellerstein. “Independence and port oracles for matroids, with an application to computational learning theory”. In: *Combinatorica* 16.2 (1996), pp. 189–208.
- [Dec+18] Wolfram Decker et al. *SINGULAR — A computer algebra system for polynomial computations*. Version 4-1-1. 2018.
- [Dor11] Dzmity Doryn. “On one example and one counterexample in counting rational points on graph hypersurfaces”. In: *Lett. Math. Phys.* 97.3 (2011), pp. 303–315.

- [Eis95] David Eisenbud. *Commutative algebra*. Vol. 150. Graduate Texts in Mathematics. With a view toward algebraic geometry. New York: Springer-Verlag, 1995, pp. xvi+785.
- [GP08] Gert-Martin Greuel and Gerhard Pfister. *A Singular introduction to commutative algebra*. extended. With contributions by Olaf Bachmann, Christoph Lossen and Hans Schönemann, With 1 CD-ROM (Windows, Macintosh and UNIX). Springer, Berlin, 2008, pp. xx+689.
- [Gri18] Darij Grinberg. *A symmetric bilinear form and a Plücker identity*. Nov. 6, 2018. MathOverflow: [a/314720](https://mathoverflow.net/a/314720). URL: <https://mathoverflow.net/a/314720>.
- [HC52] D. Hilbert and S. Cohn-Vossen. *Geometry and the imagination*. Translated by P. Néményi. Chelsea Publishing Company, New York, N. Y., 1952, pp. ix+357.
- [HS06] Craig Huneke and Irena Swanson. *Integral closure of ideals, rings, and modules*. Vol. 336. London Mathematical Society Lecture Note Series. Cambridge: Cambridge University Press, 2006, pp. xiv+431.
- [Kat16] Eric Katz. “Matroid theory for algebraic geometers”. In: *Nonarchimedean and tropical geometry*. Simons Symp. Springer, [Cham], 2016, pp. 435–517.
- [Kut74] Ronald E. Kutz. “Cohen-Macaulay rings and ideal theory in rings of invariants of algebraic groups”. In: *Trans. Amer. Math. Soc.* 194 (1974), pp. 115–129.
- [Mar10] Matilde Marcolli. *Feynman motives*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2010, pp. xiv+220.
- [Mat89] Hideyuki Matsumura. *Commutative ring theory*. Second Edition. Vol. 8. Cambridge Studies in Advanced Mathematics. Translated from the Japanese by M. Reid. Cambridge University Press, Cambridge, 1989, pp. xiv+320.
- [OT92] Peter Orlik and Hiroaki Terao. *Arrangements of hyperplanes*. Vol. 300. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Berlin: Springer-Verlag, 1992, pp. xviii+325.
- [Oxl11] James Oxley. *Matroid theory*. Second Edition. Vol. 21. Oxford Graduate Texts in Mathematics. Oxford University Press, Oxford, 2011, pp. xiv+684.
- [Pat10] Eric Patterson. “On the singular structure of graph hypersurfaces”. In: *Commun. Number Theory Phys.* 4.4 (2010), pp. 659–708.
- [Piq19] Matthieu Piquerez. *A multidimensional generalization of Symanzik polynomials*. 2019. arXiv: [1901.09797](https://arxiv.org/abs/1901.09797).
- [Sey80] P. D. Seymour. “Decomposition of regular matroids”. In: *J. Combin. Theory Ser. B* 28.3 (1980), pp. 305–359.

[Tru92] K. Truemper. *Matroid decomposition*. Academic Press, Inc., Boston, MA, 1992, pp. x+398.

G. DENHAM, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WESTERN ONTARIO, LONDON, ON, CANADA N6A 5B7

Email address: gdenham@uwo.ca

M. SCHULZE, DEPARTMENT OF MATHEMATICS, TU KAISERSLAUTERN, 67663 KAISERSLAUTERN, GERMANY

Email address: mschulze@mathematik.uni-kl.de

U. WALTHER, DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, IN 47907, USA

Email address: walther@math.purdue.edu